

**GENES IDENTIFIED AS REQUIRED FOR PROLIFERATION IN  
*ESCHERICHIA COLI***

RELATED APPLICATIONS

5           This application claims priority from U.S. Provisional Patent Application Serial Number 60/117,405 filed January 27, 1999, the disclosure of which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

10           Since the discovery of penicillin, the use of antibiotics to treat the ravages of bacterial infections has saved millions of lives. With the advent of these "miracle drugs," for a time it was popularly believed that humanity might, once and for all, be saved from the scourge of bacterial infections. In fact, during the 1980s and early 1990s, many large pharmaceutical companies cut back or eliminated antibiotics research and development. They believed that infectious disease caused by bacteria finally had been conquered and  
15           that markets for new drugs were limited. Unfortunately, this belief was overly optimistic.

          The tide is beginning to turn in favor of the bacteria as reports of drug resistant bacteria become more frequent. The United States Centers for Disease Control announced that one of the most powerful known antibiotics, vancomycin, was unable to treat an infection of the common *Staphylococcus aureus* (staph). This organism is commonly  
20           found in our environment and is responsible for many nosocomial infections. The import of this announcement becomes clear when one considers that vancomycin was used for years to treat infections caused by stubborn strains of bacteria, like staph. In short, the bacteria are becoming resistant to our most powerful antibiotics. If this trend continues, it is conceivable that we will return to a time when what are presently  
25           considered minor bacterial infections are fatal diseases.

          There are a number of causes for the predicament in which practitioners of medical arts find themselves. Over-prescription and improper prescription habits by some physicians have caused an indiscriminate increase in the availability of antibiotics to the public. The patient is also partly responsible, for even in instances where an antibiotic is  
30           the appropriate treatment, patients will often improperly use the drug, the result being yet another population of bacteria that is resistant, in whole or in part, to traditional antibiotics.

The bacterial scourges that have haunted humanity remain, in spite of the development of modern scientific practices to deal with the diseases that they cause. Drug resistant bacteria are now advancing on the health of humanity. A new generation of antibiotics to once again deal with the pending health threat that bacteria present is required.

#### Discovery of New Antibiotics

As more and more bacterial strains become resistant to the panel of available antibiotics, new compounds are required. In the past, practitioners of pharmacology would have to rely upon traditional methods of drug discovery to generate novel, safe and efficacious compounds for the treatment of disease. Traditional drug discovery methods involve blindly testing potential drug candidate-molecules, often selected at random, in the hope that one might prove to be an effective treatment for some disease. The process is painstaking and laborious, with no guarantee of success. Today, the average cost to discover and develop a new drug is nearly US \$500 million, and the average time is 15 years from laboratory to patient. Improving this process, even incrementally, would represent a huge advance in the generation of novel antimicrobial agents.

Newly emerging practices in drug discovery utilize a number of biochemical techniques to provide for directed approaches to creating new drugs, rather than discovering them at random. For example, gene sequences and proteins encoded thereby that are required for the proliferation of an organism make for excellent targets since exposure of bacteria to compounds active against these targets would result in the inactivation of the organism. Once a target is identified, biochemical analysis of that target can be used to discover or to design molecules that interact with and alter the functions of the target. Using physical and computational techniques, to analyze structural and biochemical targets in order to derive compounds that interact with a target is called rational drug design and offers great future potential. Thus, emerging drug discovery practices use molecular modeling techniques, combinatorial chemistry approaches, and other means to produce and screen and/or design large numbers of candidate compounds.

Nevertheless, while this approach to drug discovery is clearly the way of the future, problems remain. For example, the initial step of identifying molecular targets

for investigation can be an extremely time consuming task. It may also be difficult to design molecules that interact with the target by using computer modeling techniques. Furthermore, in cases where the function of the target is not known or is poorly understood, it may be difficult to design assays to detect molecules that interact with and alter the functions of the target. To improve the rate of novel drug discovery and development, methods of identifying important molecular targets in pathogenic microorganisms and methods for identifying molecules that interact with and alter the functions of such molecular targets are urgently required.

*Escherichia coli* represents an excellent model system to understand bacterial biochemistry and physiology. The estimated 4288 genes scattered along the  $4.6 \times 10^6$  base pairs of the *Escherichia coli* (*E. coli*) chromosome offer tremendous promise for the understanding of bacterial biochemical processes. In turn, this knowledge will assist in the development of new tools for the diagnosis and treatment of bacteria-caused human disease. The entire *E. coli* genome has been sequenced, and this body of information holds a tremendous potential for application to the discovery and development of new antibiotic compounds. Yet, in spite of this accomplishment, the general functions or roles of many of these genes are still unknown. For example, the total number of proliferation-required genes contained within the *E. coli* genome is unknown, but has been variously estimated at around 200 to 700 (Armstrong, K.A. and Fan, D.P. Essential Genes in the *metB-malB* Region of *Escherichia coli* K12, 1975, J. Bacteriol. 126: 48-55).

Novel, safe and effective antimicrobial compounds are needed in view of the rapid rise of antibiotic resistant microorganisms. However, prior to this invention, the characterization of even a single bacterial gene was a painstaking process, requiring years of effort. Accordingly, there is an urgent need for more novel methods to identify and characterize bacterial genomic sequences that encode gene products required for proliferation and for methods to identify molecules that interact with and alter the functions of such genes and gene products.

#### SUMMARY OF THE INVENTION

One embodiment of the present invention is a purified or isolated nucleic acid sequence consisting essentially of one of SEQ ID NOs: 1-81, 405-485, wherein said

nucleic acid inhibits microorganism proliferation. The nucleic acid sequence may be complementary to at least a portion of a coding sequence of a gene whose expression is required for microorganism proliferation. The nucleic acid sequence may comprise a fragment of one of SEQ ID NOs. 1-81, 405-485, said fragment selected from the group consisting of fragments comprising at least 10, at least 20, at least 25, at least 30, at least 50 or more than 50 consecutive bases of one of SEQ ID NOs: 1-81, 405-485. The nucleic acid sequence may be complementary to a coding sequence of a gene whose expression is required for microorganism proliferation.

Another embodiment of the present invention is a vector comprising a promoter operably linked to a nucleic acid comprising a sequence selected from the group consisting of SEQ ID NOs. 1-81, 405-485. The promoter may be active in an organism selected from the group consisting of *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Helicobacter pylori*, *Neisseria gonorrhoeae*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Salmonella typhimurium*, *Saccharomyces cerevisiae*, *Candida albicans*, *Cryptococcus neoformans*, *Aspergillus fumigatus*, *Klebsiella pneumoniae*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella choleraesuis*, *Staphylococcus epidermidis*, *Mycobacterium tuberculosis*, *Mycobacterium leprae*, *Treponema pallidum*, *Bacillus anthracis*, *Yersinia pestis*, *Clostridium botulinum*, *campylobacter jejuni*, *Chlamydia trachomatis*, *Chlamydia pneumoniae* or any species falling within the genera of any of the above species.

Another embodiment of the present invention is a host cell containing the vectors described above.

Another embodiment of the present invention is a purified or isolated nucleic acid consisting essentially of the coding sequence of one of SEQ ID NOs: 82-88, 90-242. One aspect of this embodiment is a fragment of the nucleic acid comprising at least 10, at least 20, at least 25, at least 30, at least 50 or more than 50 consecutive bases of one of SEQ ID NOs: 82-88, 90-242.

Another embodiment of the present invention is a vector comprising a promoter operably linked to the nucleic acids of the preceding embodiment.

Another aspect of the present invention is a purified or isolated nucleic acid comprising a nucleic acid sequence complementary to at least a portion of an intragenic



sequence, intergenic sequence, sequences spanning at least a portion of two or more genes, 5' noncoding region, or 3' noncoding region within an operon encoding a polypeptide comprising a sequence selected from the group consisting of SEQ ID NOs: 243-357, 359-398.

5 Another embodiment of the present invention is a purified or isolated nucleic acid comprising a nucleic acid having at least 70% homology to a sequence selected from the group consisting of SEQ ID NOs 1-81, 405-485, 82-88, 90-242 or the sequences complementary thereto as determined using BLASTN version 2.0 with the default parameters. The nucleic acid may be from an organism selected from the group  
10 consisting of *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Helicobacter pylori*, *Neisseria gonorrhoeae*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Salmonella typhimurium*, *Saccharomyces cerevisiae*, *Candida albicans*, *Cryptococcus neoformans*, *Aspergillus fumigatus*, *Klebsiella pneumoniae*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella choleraesuis*, *Staphylococcus epidermidis*, *Mycobacterium tuberculosis*, *Mycobacterium leprae*, *Treponema pallidum*, *Bacillus anthracis*, *Yersinia pestis*, *Clostridium botulinum*, *campylobacter jejuni*, and *Chlamydia trachomatis*, *Chlamydia pneumoniae*  
15 or any species falling within the genera of any of the above species.

Another embodiment of the present invention is a purified or isolated nucleic acid consisting essentially of a nucleic acid encoding a polypeptide having a sequence  
20 selected from the group consisting of SEQ ID NOs.: 243-357, 359-398.

Another embodiment of the present invention is a vector comprising a promoter operably linked to a nucleic acid encoding a polypeptide having a sequence selected from the group consisting of SEQ ID NOs.: 243-357, 359-398.

25 Another embodiment of the present invention is a host cell containing the vector of the preceding embodiment.

Another embodiment of the present invention is purified or isolated polypeptide comprising the sequence of one of SEQ ID NOs: 243-357, 359-398.

Another embodiment of the present invention is purified or isolated polypeptide  
30 comprising a fragment of one of the polypeptides of SEQ ID NOs. 243-357, 359-398, said fragment selected from the group consisting of fragments comprising at least 5, at

least 10, at least 20, at least 30, at least 40, at least 50, at least 60 or more than 60 consecutive amino acids of one of the polypeptides of SEQ ID NOs.: 243-357, 359-398.

Another embodiment of the present invention is an antibody capable of specifically binding the polypeptide of the preceding embodiment.

5 Another embodiment of the present invention is method of producing a polypeptide, comprising introducing a vector comprising a promoter operably linked to a nucleic acid encoding a polypeptide having a sequence selected from the group consisting of SEQ ID NOs. 243-357, 359-398 into a cell. The method may further comprise the step of isolating said protein.

10 Another embodiment of the present invention is a method of inhibiting proliferation comprising inhibiting the activity or reducing the amount of a polypeptide having a sequence selected from the group consisting of SEQ ID NOs. 243-357, 359-398 or inhibiting the activity or reducing the amount of a nucleic acid encoding said polypeptide.

15 Another embodiment of the present invention is method for identifying compounds which influence the activity of a polypeptide required for proliferation comprising:

contacting a polypeptide comprising a sequence selected from the group consisting of 243-357, 359-398 with a candidate compound; and

20 determining whether said compound influences the activity of said polypeptide.

The activity may be an enzymatic activity. The activity may be a carbon compound catabolism activity. The activity may be a biosynthetic activity. The activity may be a transporter activity. The activity may be a transcriptional activity. The activity may be a DNA replication activity. The activity may be a cell division activity.

25 Another embodiment of the present invention is a compound identified using the above method.

Another embodiment of the present invention is method for assaying compounds for the ability to reduce the activity or level of a polypeptide required for proliferation, comprising:

30 providing a target, wherein said target comprises the coding sequence of a sequence selected from the group consisting of SEQ ID NOs. 82-88, 90-242;

contacting said target with a candidate compound; and

measuring an activity of said target.

The target may be a messenger RNA molecule transcribed from a coding region of one of SEQ ID. NOs.: 82-88, 90-242 and said activity is translation of said messenger RNA. The target may be a coding region of one of SEQ ID. NOs. 82-88, 90-242 and said activity is transcription of said messenger RNA.

Another embodiment of the present invention is a compound identified using the method above.

Another embodiment of the present invention is a method for identifying compounds which reduce the activity or level of a gene product required for cell proliferation comprising the steps of:

expressing an antisense nucleic acid against a nucleic acid encoding said gene product in a cell to reduce the activity or amount of said gene product in said cell, thereby producing a sensitized cell;

contacting said sensitized cell with a compound; and

determining whether said compound inhibits the growth of said sensitized cell to a greater extent than said compound inhibits the growth of a nonsensitized cell.

The cell may be selected from the group consisting of bacterial cells, fungal cells, plant cells, and animal cells. The cell may be an *E. coli* cell. The cell may be from an organism selected from the group consisting of *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Helicobacter pylori*, *Neisseria gonorrhoeae*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Salmonella typhimurium*, *Saccharomyces cerevisiae*, *Candida albicans*, *Cryptococcus neoformans*, *Aspergillus fumigatus*, *Klebsiella pneumoniae*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella choleraesuis*, *Staphylococcus epidermidis*, *Mycobacterium tuberculosis*, *Mycobacterium leprae*, *Treponema pallidum*, *Bacillus anthracis*, *Yersinia pestis*, *Clostridium botulinum*, *Campylobacter jejuni*, and *Chlamydia trachomatis*, *Chlamydia pneumoniae* or any species falling within the genera of any of the above species. The antisense nucleic acid may be transcribed from an inducible promoter. The method may, further comprise the step of contacting said cell with a concentration of inducer which induces said antisense nucleic acid to a sublethal level. The sub-lethal concentration of said inducer may be such that growth inhibition is 8% or more. The inducer may be isopropyl-1-thio- $\beta$ -D-galactoside. The growth inhibition

may be measured by monitoring optical density of a culture growth solution. The gene product may be a polypeptide. The gene product may be an RNA. The gene product may comprise a polypeptide having a sequence selected from the group consisting of SEQ ID NOs.: 243-357, 359-398.

5           Another embodiment of the present invention is a compound identified using the method above.

          Another embodiment of the present invention is a method for inhibiting cellular proliferation comprising introducing a compound with activity against a gene corresponding to one of SEQ ID NOs.: 82-88, 90-242 or with activity against the  
10   product of said gene into a population of cells expressing a gene. The compound may be an antisense oligonucleotide comprising a sequence selected from the group consisting of SEQ ID NOs.: 1-81, 405-485, or a proliferation-inhibiting portion thereof. The proliferation inhibiting portion of one of SEQ ID NOs. 1-81, 405-485 may be a fragment comprising at least 10, at least 20, at least 25, at least 30, at least 50 or more  
15   than 50 consecutive bases of one of SEQ ID NOs: 1-81, 405-485. The compound may be a triple helix oligonucleotide.

          Another embodiment of the present invention is a preparation comprising an effective concentration of an antisense oligonucleotide comprising a sequence selected from the group consisting of SEQ ID NOs.: 1-81, 405-485, or a proliferation-inhibiting  
20   portion thereof in a pharmaceutically acceptable carrier. The proliferation-inhibiting portion of one of SEQ ID NOs. 1-81, 405-485 may comprise at least 10, at least 20, at least 25, at least 30, at least 50 or more than 50 consecutive bases of one of SEQ ID NOs: 1-81, 405-485.

          Another embodiment of the present invention is a method for inhibiting the  
25   expression of a gene in an operon required for proliferation comprising contacting a cell in a cell population with an antisense nucleic acid, said cell expressing a gene corresponding to one of SEQ ID NOs.: 82-88, 90-242, wherein said antisense nucleic acid comprises at least a proliferation-inhibiting portion of said operon in an antisense orientation that is effective in inhibiting expression of said gene. The antisense nucleic  
30   acid may be complementary to a sequence of a gene comprising one or more of SEQ ID NOs.: 82-88, 90-242. The antisense nucleic acid may be a sequence of one of SEQ ID NOs.: 1-81, 405-485, or a portion thereof. The cell may be contacted with said

antisense nucleic acid by introducing a plasmid which expresses said antisense nucleic acid into said cell population. The cell may be contacted with said antisense nucleic acid by introducing a phage which expresses said antisense nucleic acid into said cell population. The cell may be contacted with said antisense nucleic acid by introducing a  
5 sequence encoding said antisense nucleic acid into the chromosome of said cell into said cell population. The cell may be contacted with said antisense nucleic acid by introducing a retron which expresses said antisense nucleic acid into said cell population. The cell may be contacted with said antisense nucleic acid by introducing a ribozyme into said cell-population, wherein a binding portion of said ribozyme is  
10 complementary to said antisense oligonucleotide. The cell may be contacted with said antisense nucleic acid by introducing a liposome comprising said antisense oligonucleotide into said cell. The cell may be contacted with said antisense nucleic acid by electroporation. The antisense nucleic acid may be a fragment comprising at least 10, at least 20, at least 25, at least 30, at least 50 or more than 50 consecutive bases  
15 of one of SEQ ID NOs: 82-88, 90-242. The antisense nucleic acid may be an oligonucleotide.

Another embodiment of the present invention is a method for identifying bacterial strains comprising the steps of:

providing a sample containing a bacterial species; and  
20 identifying a bacterial species using a species specific probe having a sequence selected from the group consisting of SEQ ID NOs. 1-81, 405-485, 82-88, 90-242.

Another embodiment of the present invention is a method for identifying a gene in a microorganism required for proliferation comprising:

- (a) identifying an inhibitory nucleic acid which inhibits the activity of a gene or  
25 gene product required for proliferation in a first microorganism;  
(b) contacting a second microorganism with said inhibitory nucleic acid;  
(c) determining whether said inhibitory nucleic acid from said first microorganism inhibits proliferation of said second microorganism; and  
(d) identifying the gene in said second microorganism which is inhibited by said  
30 inhibitory nucleic acid.

Another embodiment of the present invention is a method for assaying a compound for the ability to inhibit proliferation of a microorganism comprising:

- (a) identifying a gene or gene product required for proliferation in a first microorganism;
- (b) identifying a homolog of said gene or gene product in a second microorganism;
- (c) identifying an inhibitory nucleic acid sequence which inhibits the activity of said homolog in said second microorganism;
- (d) contacting said second microorganism with a proliferation-inhibiting amount of said inhibitory nucleic acid, thus sensitizing said second microorganism;
- (e) contacting the sensitized microorganism of step (d) with a compound; and
- (f) determining whether said compound inhibits proliferation of said sensitized microorganism to a greater extent than said compound inhibits proliferation of a nonsensitized microorganism.

The step of identifying a gene involved in proliferation in a first microorganism may comprise:

introducing a nucleic acid comprising a random genomic fragment from said first microorganism operably linked to a promoter wherein said random genomic fragment is in the antisense orientation; and

comparing the proliferation of said first microorganism transcribing a first level of said random genomic fragment to the proliferation of said first microorganism transcribing a lower level of said random genomic fragment, wherein a difference in proliferation indicates that said random genomic fragment comprises a gene involved in proliferation.

The step of identifying a homolog of said gene in a second microorganism may comprise identifying a homologous nucleic acid or a nucleic acid encoding a homologous polypeptide in a database using an algorithm selected from the group consisting of BLASTN version 2.0 with the default parameters and FASTA version 3.0t78 algorithm with the default parameters. The step of identifying a homolog of said gene in a second microorganism may comprise identifying a homologous nucleic acid or a nucleic acid encoding a homologous polypeptide by identifying nucleic acids which hybridize to said first gene. The step of identifying a homolog of said gene in a second microorganism may comprise expressing a nucleic acid which inhibits the proliferation of said first microorganism in said second microorganism. The inhibitory nucleic acid may be an antisense nucleic acid. The inhibitory nucleic acid may comprise an

antisense nucleic acid to a portion of said homolog. The inhibitory nucleic acid may comprise an antisense nucleic acid to a portion of the operon encoding said homolog. The step of contacting the second microorganism with a proliferation-inhibiting amount of said nucleic acid sequence may comprise directly contacting said second  
5 microorganism with said nucleic acid. The step of contacting the second microorganism with a proliferation-inhibiting amount of said nucleic acid sequence may comprise expressing an antisense nucleic acid to said homolog in said second microorganism.

Another embodiment of the present invention is a compound identified using the  
10 method above.

Another embodiment of the present invention is a method of assaying a compound for the ability to inhibit proliferation comprising:

- (a) identifying an inhibitory nucleic acid sequence which inhibits the activity of a gene or gene product required for proliferation in a first microorganism;
- 15 (b) contacting a second microorganism with a proliferation-inhibiting amount of said inhibitory nucleic acid, thus sensitizing said second microorganism;
- (c) contacting the proliferation-inhibited microorganism of step (b) with a compound; and
- (d) determining whether said compound inhibits proliferation of said sensitized  
20 second microorganism to a greater extent than said compound inhibits proliferation of a nonsensitized second microorganism.

The inhibitory nucleic acid may be an antisense nucleic acid which inhibits the proliferation of said first microorganism. The inhibitory nucleic acid may comprise a portion of an antisense nucleic acid which inhibits the proliferation of said first  
25 microorganism. The inhibitory nucleic acid may comprise an antisense molecule against the entire coding region of the gene involved in proliferation of the first microorganism. The inhibitory nucleic acid may comprise an antisense nucleic acid to a portion of the operon encoding the gene involved in proliferation of the first microorganism.

Another embodiment of the present invention is a compound identified using the  
30 method above.

Another embodiment of the present invention is a method for assaying compounds for activity against a biological pathway required for proliferation comprising:

sensitizing a cell by expressing an antisense nucleic acid against a nucleic acid encoding a gene product required for proliferation in a cell to reduce the activity or amount of said gene product;

contacting the sensitized cell with a compound; and

determining whether said compound inhibits the growth of said sensitized cell to a greater extent than said compound inhibits the growth of a nonsensitized cell.

The cell may be selected from the group consisting of bacterial cells, fungal cells, plant cells, and animal cells. The cell may be an *E. coli* cell. The cell may be an organism selected from the group consisting of *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Helicobacter pylori*, *Neisseria gonorrhoeae*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Salmonella typhimurium*, *Saccharomyces cerevisiae*, *Candida albicans*, *Cryptococcus neoformans*, *Aspergillus fumigatus*, *Klebsiella pneumoniae*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella choleraesuis*, *Staphylococcus epidermidis*, *Mycobacterium tuberculosis*, *Mycobacterium leprae*, *Treponema pallidum*, *Bacillus anthracis*, *Yersinia pestis*, *Clostridium botulinum*, *Campylobacter jejuni*, and *Chlamydia trachomatis*, *Chlamydia pneumoniae* or any species falling within the genera of any of the above species. The antisense nucleic acid may be transcribed from an inducible promoter. The method may further comprise contacting the cell with an agent which induces expression of said antisense nucleic acid from said inducible promoter, wherein said antisense nucleic acid is expressed at a sublethal level. The sublethal level of said antisense nucleic acid may inhibit proliferation by 8% or more. The agent may be isopropyl-1-thio- $\beta$ -D-galactoside (IPTG). The inhibition of proliferation may be measured by monitoring the optical density of a liquid culture. The gene product may comprise a polypeptide having a sequence selected from the group consisting of SEQ ID NOs: 243-357, 359-398.

Another embodiment of the present invention is a compound identified using the method above.



Another embodiment of the present invention is a method for assaying a compound for the ability to inhibit cellular proliferation comprising:

contacting a cell with an agent which reduces the activity or level of a gene product required for proliferation of said cell;

5           contacting said cell with said compound; and

determining whether said compound reduces proliferation to a greater extent than said compound reduces proliferation of cells which have not been contacted with said agent.

10           The agent which reduces the activity or level of a gene product required for proliferation of said cell may comprise an antisense nucleic acid to a gene or operon required for proliferation. The agent which reduces the activity or level of a gene product required for proliferation of said cell may comprise an antibiotic. The cell may contain a temperature sensitive mutation which reduces the activity or level of said gene product required for proliferation of said cell. The antisense nucleic acid may be  
15           directed against the same functional domain of said gene product required for proliferation of said cell to which said antisense nucleic acid is directed. The antisense nucleic acid may be directed against a different functional domain of said gene product required for proliferation of said cell than the functional domain to which said antisense nucleic acid is directed.

20           Another embodiment of the present invention is a compound identified using the method above.

Another embodiment of the present invention is a method for identifying the pathway in which a proliferation-required nucleic acid or its gene product lies comprising:

25           expressing a sublethal level of an antisense nucleic acid directed against said proliferation-required nucleic acid in a cell;

contacting said cell with an antibiotic, wherein the a biological pathway on which said antibiotic acts is known; and

30           determining whether said cell has a substantially greater sensitivity to said antibiotic than a cell which does not express said sublethal level of said antisense nucleic acid.

Another embodiment of the present invention is a method for determining the pathway on which a test compound acts comprising:

(a) expressing a sublethal level of an antisense nucleic acid directed against a proliferation-required nucleic acid in a cell, wherein the biological pathway in which said proliferation-required nucleic acid lies is known,

(b) contacting said cell with said test compound; and

(c) determining whether said cell has a substantially greater sensitivity to said test compound than a cell which does not express said sublethal level of said antisense nucleic acid.

The method may further comprise:

(d) expressing a sublethal level of a second antisense nucleic acid directed against a second proliferation-required nucleic acid in said cell, wherein said second proliferation-required nucleic acid is in a different biological pathway than said proliferation-required nucleic acid in step (a); and

(e) determining whether said cell has a substantially greater sensitivity to said test compound than a cell which does not express said sublethal level of said second antisense nucleic acid.

Another embodiment of the present invention is a purified or isolated nucleic acid consisting essentially of one of SEQ ID NOs: 358, 399-402.

Another embodiment of the present invention is a purified or isolated nucleic acid comprising a sequence selected from the group consisting of 1-81, 405-485, 82-88, 90-242, 358, 399-402.

Another embodiment of the present invention is a compound which interacts with the gene or gene product of a nucleic acid comprising a sequence of one of SEQ ID NOs: 82-88, 90-242 to inhibit proliferation.

Another embodiment of the present invention compound which interacts with a polypeptide comprising one of SEQ ID NOs. 243-357, 359-398 to inhibit proliferation.

Another embodiment of the present invention is a compound which interacts with a nucleic acid comprising one of SEQ ID NOs: 358, 399-402 to inhibit proliferation.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is an IPTG dose response curve in *E. coli* transformed with an IPTG-inducible plasmid containing either an antisense clone to the *E. coli* ribosomal protein rplW (AS-rplW) which is required for protein synthesis and essential cell proliferation, or an antisense clone to the *elaD* (AS-*elaD*) gene which is not known to be involved in protein synthesis and which is also essential for proliferation.

Figure 2A is a tetracycline dose response curve in *E. coli* transformed with an IPTG-inducible plasmid containing antisense to rplW(AS-rplW) in the presence of 0, 20 or 50  $\mu$ M IPTG.

Figure 2B is a tetracycline dose response curve in *E. coli* transformed with an IPTG-inducible plasmid containing antisense to *elaD* (AS-*elaD*) in the presence of 0, 20 or 50  $\mu$ M IPTG.

Figure 3 is a graph showing the fold increase in tetracycline sensitivity of *E. coli* transfected with antisense clones to essential ribosomal proteins L23 (AS-rplW) and L7/L12 and L10 (AS-rplLrplJ). Antisense clones to genes known not to be involved in protein synthesis (atpB/E(AS-atpB/E ), visC (AS-visC, *elaD* (AS-*elaD*), yohH (AS-yohH) are much less sensitive to tetracycline.

#### Definitions

By "biological pathway" is meant any discrete cell function or process that is carried out by a gene product or a subset of gene products. Biological pathways include enzymatic, biochemical and metabolic pathways as well as pathways involved in the production of cellular structures such cell walls. Biological pathways that are usually required for proliferation of microorganisms include, but are not limited to, cell division, DNA synthesis & replication, RNA synthesis (transcription), protein synthesis (translation), protein processing, protein transport, fatty acid biosynthesis, cell wall synthesis, cell membrane synthesis & maintenance, etc.

By "inhibit activity against a gene or gene product" is meant having the ability to interfere with the function of a gene or gene product in such a way as to decrease expression of the gene or to reduce the level or activity of a product of the gene. Agents which have activity against a gene include agents that inhibit transcription of the gene and agents that inhibit translation of the mRNA transcribed from the gene. In microorganisms, agents which have activity against a gene can act to decrease expression of the operon in which the gene resides or alter the processing of operon

RNA such as to reduce the level or activity of the gene product. The gene product can be a non-translated RNA such as ribosomal RNA, a translated RNA (mRNA) or the protein product resulting from translation of the gene mRNA. Of particular utility to the present invention are anti-sense RNAs that have activities against the operons or genes to which they specifically hybridize.

By "activity against a gene product" is meant having the ability to inhibit the function or to reduce the level or activity of the gene product in a cell.

By "activity against a protein" is meant having the ability to inhibit the function or to reduce the level or activity of the protein in a cell.

By "activity against nucleic acid" is meant having the ability to inhibit the function or to reduce the level or activity of the nucleic acid in a cell.

As used herein, "sublethal" means a concentration of an agent below the concentration required to inhibit all cell growth.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention describes a group of *E. coli* genes and gene families required for growth and/or proliferation. A proliferation-required gene or gene family is one where, in the absence of a gene transcript and/or gene product, growth or viability of the microorganism is reduced or eliminated. Thus, as used herein the terminology "proliferation-required" or "required for proliferation" encompasses sequences where the absence of a gene transcript and/or gene product completely eliminates cell growth as well as sequences where the absence of a gene transcript and/or gene product merely reduces cell growth. These proliferation-required genes can be used as potential targets for the generation of new antimicrobial agents. To achieve that goal, the present invention also encompasses novel assays for analyzing proliferation-required genes and for identifying compounds which interact with the gene products of the proliferation-required genes. In addition, the present invention contemplates the expression of genes and the purification of the proteins encoded by the nucleic acid sequences identified as required proliferation genes and reported herein. The purified proteins can be used to generate reagents and screen small molecule libraries or other candidate compound libraries for compounds that can be further developed to yield novel antimicrobial compounds. The present invention also describes methods for identification of homologous genes in organisms other than *E. coli*.

The present invention utilizes a novel method to identify proliferation-required *E. coli* sequences. Generally, a library of nucleic acid sequences from a given source are subcloned or otherwise inserted into an inducible expression vector, thus forming an expression library. Although the insert nucleic acids may be derived from the chromosome of the organism into which the expression vector is to be introduced, because the insert is not in its natural chromosomal location, the insert nucleic acid is an exogenous nucleic acid for the purposes of the discussion herein. The term expression is defined as the production of an RNA molecule from a gene, gene fragment, genomic fragment, or operon. Expression can also be used to refer to the process of peptide or polypeptide synthesis. An expression vector is defined as a vehicle by which a ribonucleic acid (RNA) sequence is transcribed from a nucleic acid sequence carried within the expression vehicle. The expression vector can also contain features that permit translation of a protein product from the transcribed RNA message expressed from the exogenous nucleic acid sequence carried by the expression vector. Accordingly, an expression vector can produce an RNA molecule as its sole product or the expression vector can produce a RNA molecule that is ultimately translated into a protein product.

Once generated, the expression library containing the exogenous nucleic acid sequences is introduced into an *E. coli* population to search for genes that are required for bacterial proliferation. Because the library molecules are foreign to the population of *E. coli*, the expression vectors and the nucleic acid segments contained therein are considered exogenous nucleic acid.

Expression of the exogenous nucleic acid fragments in the test population of *E. coli* containing the expression vector library is then activated. Activation of the expression vectors consists of subjecting the cells containing the vectors to conditions that result in the expression of the exogenous nucleic acid sequences carried by the expression vector library. The test population of *E. coli* cells is then assayed to determine the effect of expressing the exogenous nucleic acid fragments on the test population of cells. Those expression vectors that, upon activation and expression, negatively impact the growth of the *E. coli* screen population were identified, isolated, and purified for further study.

A variety of assays are contemplated to identify nucleic acid sequences that negatively impact growth upon expression. In one embodiment, growth in *E. coli* cultures expressing exogenous nucleic acid sequences and growth in cultures not expressing these

sequences is compared. Growth measurements are assayed by examining the extent of growth by measuring optical densities. Alternatively, enzymatic assays can be used to measure bacterial growth rates to identify exogenous nucleic acid sequences of interest. Colony size, colony morphology, and cell morphology are additional factors used to evaluate growth of the host cells. Those cultures that failed to grow or grow with reduced efficiency under expression conditions are identified as containing an expression vector encoding a nucleic acid fragment that negatively affects a proliferation-required gene.

Once exogenous nucleic acid sequences of interest are identified, they are analyzed. The first step of the analysis is to acquire the nucleic acid sequence of the nucleic acid fragment of interest. To achieve this end, the insert in those expression vectors identified as containing a sequence of interest is sequenced, using standard techniques well known in the art. The next step of the process is to determine the source of the nucleic acid sequence.

Determination of sequence source is achieved by comparing the obtained sequence data with known sequences in various genetic databases. The sequences identified are used to probe these gene databases. The result of this procedure is a list of exogenous nucleic acid sequences corresponding to a list that includes novel bacterial genes required for proliferation as well as genes previously identified as required for proliferation.

The number of DNA and protein sequences available in database systems has been growing exponentially for years. For example, at the end of 1998, the complete sequences of *Caenorhabditis elegans*, *Saccharomyces cerevisiae* and nineteen bacterial genomes, including *E. coli* were available. This sequence information is stored in a number of databanks, such as GenBank (the National Center for Biotechnology Information (NCBI), and is publicly available for searching.

A variety of computer programs are available to assist in the analysis of the sequences stored within these databases. FastA, (W. R. Pearson (1990) "Rapid and Sensitive Sequence Comparison with FASTP and FASTA" Methods in Enzymology 183:63- 98), Sequence Retrieval System (SRS), (Etzold & Argos, SRS an indexing and retrieval tool for flat file data libraries. Comput. Appl. Biosci. 9:49-57, 1993) are two examples of computer programs that can be used to analyze sequences of interest. In one embodiment of the present invention, the BLAST family of computer programs,

which includes BLASTN version 2.0 with the default parameters, or BLASTX version 2.0 with the default parameters, is used to analyze nucleic acid sequences.

BLAST, an acronym for "Basic Local Alignment Search Tool," is a family of programs for database similarity searching. The BLAST family of programs includes: BLASTN, a nucleotide sequence database searching program, BLASTX, a protein database searching program where the input is a nucleic acid sequence; and BLASTP, a protein database searching program. BLAST programs embody a fast algorithm for sequence matching, rigorous statistical methods for judging the significance of matches, and various options for tailoring the program for special situations. Assistance in using the program can be obtained by e-mail at [blast@ncbi.nlm.nih.gov](mailto:blast@ncbi.nlm.nih.gov).

Bacterial genes are often transcribed in polycistronic groups. These groups comprise operons, which are a collection of genes and intergenic sequences. The genes of an operon are co-transcribed and are often related functionally. Given the nature of the screening protocol, it is possible that the identified exogenous nucleic acid sequence corresponds to a gene or portion thereof with or without adjacent noncoding sequences, an intragenic sequence (i.e. a sequence within a gene), an intergenic sequence (i.e. a sequence between genes), a sequence spanning at least a portion of two or more genes, a 5' noncoding region or a 3' noncoding region located upstream or downstream from the actual sequence that is required for bacterial proliferation. Accordingly, determining which of the genes that are encoded within the operons are individually required for proliferation is often desirable.

In one embodiment of the present invention, an operon is dissected to determine which gene or genes are required for proliferation. For example, the RegulonDB DataBase described by Huerta et al. (*Nucl. Acids Res.* **26**:55-59, 1998), which may also be found on the website [http://www.cifn.unam.mx/Computational\\_Biology/regulondb/](http://www.cifn.unam.mx/Computational_Biology/regulondb/), may be used to identify the boundaries of operons encoded within microbial genomes. A number of techniques that are well known in the art can be used to dissect the operon. In one aspect of this embodiment, gene disruption by homologous recombination is used to individually inactivate the genes of an operon that is thought to contain a gene required for proliferation.

Several gene disruption techniques have been described for the replacement of a functional gene with a mutated, non-functional (null) allele. These techniques generally

involve the use of homologous recombination. The method described by Link et al. (J. Bacteriol 1997 179:6228; incorporated herein by reference in it's entirety) serves as an excellent example of these methods as applicable to disruption of genes in *E. coli*. This technique uses crossover PCR to create a null allele with an in-frame deletion of the coding region of a target gene. The null allele is constructed in such a way that sequences adjacent to the wild type gene (ca. 500 bp) are retained. These homologous sequences surrounding the deletion null allele provide targets for homologous recombination so that the wild type gene on the *E. coli* chromosome can be replaced by the constructed null allele.

The crossover PCR amplification product is subcloned into the vector pKO3, the features of which include a chloramphenicol resistance gene, the counter-selectable marker *sacB*, and a temperature sensitive autonomous replication function. Following transformation of an *E. coli* cell population with such a vector, selection for cells that have undergone homologous recombination of the vector into the chromosome is achieved by growth on chloramphenicol at the non-permissive temperature of 43°C. Under these conditions, autonomous replication of the plasmid cannot occur and cell are resistant to chloramphenicol only if the chloramphenicol resistance gene has been integrated into the chromosome. Usually a single crossover event is responsible for this integration event such that the *E. coli* chromosome now contains a tandem duplication of the target gene consisting of one wild type allele and one deletion null allele separated by vector sequence.

This new *E. coli* strain containing the tandem duplication can be maintained at permissive temperatures in the presence of drug selection (chloramphenicol). Subsequently, cells of this new strain are cultured at the permissive temperature 30°C without drug selection. Under these conditions, the chromosome of some of the cells within the population will have undergone an internal homologous recombination event resulting in removal of the plasmid sequences. Subsequent culturing of the strain in growth medium lacking chloramphenicol but containing sucrose is used to select for such recombinative resolutions. In the presence of the counter-selectable marker *sacB*, sucrose is rendered into a toxic metabolite. Thus, cells that survive this counter-selection have lost both the plasmid sequences from the chromosome and the



autonomously replicating plasmid that results as a byproduct of recombinative resolution.

There are two possible outcomes of the above recombinative resolution via homologous recombination. Either the wild type copy of the targeted gene is retained on the chromosome or the mutated null allele is retained on the chromosome. In the case of an essential gene, a single copy of the null allele would be lethal and such cells should not be obtained by the above procedure when applied to essential genes. In the case of a non-essential gene, roughly equal numbers of cells containing null alleles and cells containing wild type alleles should be obtained. Thus, the method serves as a test for essentiality of the targeted gene: when applied to essential genes, only cells with a wild type allele on the chromosome will be obtained.

Other techniques have also been described for the creation of disruption mutations in *E. coli*. For example, Link et al. also describe inserting an in-frame sequence tag concomitantly with an in-frame deletion in order to simplify analysis of recombinants obtained. Further, Link et al. describe disruption of genes with a drug resistance marker such as a kanamycin resistance gene. Arigoni et al., (Arigoni, F. et al. A Genome-based Approach for the Identification of Essential Bacterial Genes, Nature Biotechnology 16: 851-856, the disclosure of which is incorporated herein by reference in its entirety) describe the use of gene disruption combined with engineering a second copy of a test gene such that the expression of the gene is regulated by and inducible promoter such as the arabinose promoter to test the essentiality of the gene. Many of these techniques result in the insertion of large fragments of DNA into the gene of interest, such as a drug selection marker. An advantage of the technique described by Link et al. is that it does not rely on an insertion into the gene to cause a functional defect, but rather results in the precise removal of the coding region. This insures the lack of polar effects on the expression of genes downstream from the target gene.

Recombinant DNA techniques can be used to express the entire coding sequences of the gene identified as required for proliferation, or portions thereof. The over-expressed proteins can be used as reagents for further study. The identified exogenous sequences are isolated, purified, and cloned into a suitable expression vector using methods well known in the art. If desired, the nucleic acids can contain the sequences encoding a signal peptide to facilitate secretion of the expressed protein.

Expression of fragments of the bacterial genes identified as required for proliferation is also contemplated by the present invention. The fragments of the identified genes can encode a polypeptide comprising at least 5, at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55, at least 60, at least 65, at least 75, or more than 75 consecutive amino acids of a gene complementary to one of the identified sequences of the present invention. The nucleic acids inserted into the expression vectors can also contain sequences upstream and downstream of the coding sequence.

When expressing the coding sequence of an entire gene identified as required for bacterial proliferation or a fragment thereof, the nucleic acid sequence to be expressed is operably linked to a promoter in an expression vector using conventional cloning technology. The expression vector can be any of the bacterial, insect, yeast, or mammalian expression systems known in the art. Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, MA), Stratagene (La Jolla, California), Promega (Madison, Wisconsin), and Invitrogen (San Diego, California). If desired, to enhance expression and facilitate proper protein folding, the codon usage and codon bias of the sequence can be optimized for the particular expression organism in which the expression vector is introduced, as explained by Hatfield, et al., U.S. Patent No. 5,082,767, incorporated herein by this reference. Fusion protein expression systems are also contemplated by the present invention.

Following expression of the protein encoded by the identified exogenous nucleic acid sequence, the protein is purified. Protein purification techniques are well known in the art. Proteins encoded and expressed from identified exogenous nucleic acid sequences can be partially purified using precipitation techniques, such as precipitation with polyethylene glycol. Chromatographic methods usable with the present invention can include ion-exchange chromatography, gel filtration, use of hydroxyapatite columns, immobilized reactive dyes, chromatofocusing, and use of high-performance liquid chromatography. Electrophoretic methods such one-dimensional gel electrophoresis, high-resolution two-dimensional polyacrylamide electrophoresis, isoelectric focusing, and others are contemplated as purification methods. Also, affinity chromatographic methods, comprising antibody columns, ligand presenting columns and other affinity

chromatographic matrices are contemplated as purification methods in the present invention.

The purified proteins produced from the gene coding sequences identified as required for proliferation can be used in a variety of protocols to generate useful antimicrobial reagents. In one embodiment of the present invention, antibodies are generated against the proteins expressed from the identified exogenous nucleic acid sequences. Both monoclonal and polyclonal antibodies can be generated against the expressed proteins. Methods for generating monoclonal and polyclonal antibodies are well known in the art. Also, antibody fragment preparations prepared from the produced antibodies discussed above are contemplated.

Another application for the purified proteins of the present invention is to screen small molecule libraries for candidate compounds active against the various target proteins of the present invention. Advances in the field of combinatorial chemistry provide methods, well known in the art, to produce large numbers of candidate compounds that can have a binding, or otherwise inhibitory effect on a target protein. Accordingly, the screening of small molecule libraries for compounds with binding affinity or inhibitory activity for a target protein produced from an identified gene sequence is contemplated by the present invention.

The present invention further contemplates utility against a variety of other pathogenic organisms in addition to *E. coli*. For example, the invention has utility in identifying genes required for proliferation in prokaryotes and eukaryotes. For example, the invention has utility with protists, such as *Plasmodium* spp.; plants; animals, such as *Entamoeba* spp. and *Contracaecum* spp.; and fungi including *Candida* spp., (e.g., *Candida albicans*), *Saccharomyces cerevisiae*, *Cryptococcus neoformans*, and *Aspergillus fumigatus*. In one embodiment of the present invention, monera, specifically bacteria are probed in search of novel gene sequences required for proliferation. This embodiment is particularly important given the rise of drug resistant bacteria.

The numbers of bacterial species that are becoming resistant to existing antibiotics are growing. A partial list of these organisms includes: *Staphylococcus* spp., such as *S. aureus*; *Enterococcus* spp., such as *E. faecalis*; *Pseudomonas* spp., such as *P. aeruginosa*, *Clostridium* spp., such as *C. botulinum*, *Haemophilus* spp., such as *H. influenzae*, *Enterobacter* spp., such as *E. cloacae*, *Vibrio* spp., such as *V. cholera*;

*Moraxala* spp., such as *M. catarrhalis*; *Streptococcus* spp., such as *S. pneumoniae*,  
*Neisseria* spp., such as *N. gonorrhoeae*; *Mycoplasma* spp., such as *Mycoplasma*  
*pneumoniae*; *Salmonella typhimurium*; *Helicobacter pylori*; *Escherichia coli*; and  
*Mycobacterium tuberculosis*. The sequences identified as required for proliferation in  
5 the present invention can be used to probe these and other organisms to identify  
homologous required proliferation genes contained therein.

In one embodiment of the present invention, the nucleic acid sequences disclosed  
herein are used to screen genomic libraries generated from bacterial species of interest  
other than *E. coli*. For example, the genomic library may be from *Staphylococcus aureus*,  
10 *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Helicobacter pylori*, *Neisseria*  
*gonorrhoeae*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Haemophilus*  
*influenzae*, *Salmonella typhimurium*, *Saccharomyces cerevisiae*, *Candida albicans*,  
*Cryptococcus neoformans*, *Aspergillus fumigatus*, *Klebsiella pneumoniae*, *Salmonella*  
*typhi*, *Salmonella paratyphi*, *Salmonella choleraesuis*, *Staphylococcus epidermidis*,  
15 *Mycobacterium tuberculosis*, *Mycobacterium leprae*, *Treponema pallidum*, *Bacillus*  
*anthracis*, *Yersinia pestis*, *Clostridium botulinum*, *Campylobacter jejuni*, *Chlamydia*  
*trachomatis*, *Chlamydia pneumoniae* or any species falling within the genera of any of  
the above species. Standard molecular biology techniques are used to generate genomic  
libraries from various microorganisms. In one aspect, the libraries are generated and  
20 bound to nitrocellulose paper. The identified exogenous nucleic acid sequences of the  
present invention can then be used as probes to screen the libraries for homologous  
sequences. The homologous sequences identified can then be used as targets for the  
identification of new, antimicrobial compounds with activity against more than one  
organism.

25 For example, the preceding methods may be used to isolate nucleic acids having  
a sequence with at least 97%, at least 95%, at least 90%, at least 85%, at least 80%, or at  
least 70% identity to a nucleic acid sequence selected from the group consisting of one  
of the sequences of SEQ ID NOS. 1-81, 405-485, 82-88, 90-242, fragments comprising  
at least 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, 150, 200, 300, 400, or 500 consecutive  
30 bases thereof, and the sequences complementary thereto. Identity may be measured  
using BLASTN version 2.0 with the default parameters. (Altschul, S.F. et al. Gapped  
BLAST and PSI-BLAST: A New Generation of Protein Database Search Programs.

Nucleic Acid Res. 25: 3389-3402 (1997), the disclosure of which is incorporated herein by reference in its entirety). For example, the homologous polynucleotides may have a coding sequence which is a naturally occurring allelic variant of one of the coding sequences described herein. Such allelic variants may have a substitution, deletion or addition of one or more nucleotides when compared to the nucleic acids of SEQ ID NOs: 1-81, 405-485, 82-88, 90-242 or the sequences complementary thereto.

Additionally, the above procedures may be used to isolate nucleic acids which encode polypeptides having at least 99%, 95%, at least 90%, at least 85%, at least 80%, at least 70%, at least 60%, at least 50%, or at least 40% identity or similarity to a polypeptide having the sequence of one of SEQ ID NOs: 243-357, 359-398 or fragments comprising at least 5, 10, 15, 20, 25, 30, 35, 40, 50, 75, 100, or 150 consecutive amino acids thereof as determined using the FASTA version 3.0t78 algorithm with the default parameters. Alternatively, protein identity or similarity may be identified using BLASTP with the default parameters, BLASTX with the default parameters, or TBLASTN with the default parameters. (Alschul, S.F. et al. Gapped BLAST and PSI-BLAST: A New Generation of Protein Database Search Programs, Nucleic Acid Res. 25: 3389-3402 (1997), the disclosure of which is incorporated herein by reference in its entirety).

Alternatively, homologous nucleic acids or polypeptides may be identified by searching a database to identify sequences having a desired level of homology to a nucleic acid or polypeptide involved in proliferation or an antisense nucleic acid to a nucleic acid involved in microbial proliferation. A variety of such databases are available to those skilled in the art, including GenBank and GenSeq. In some embodiments, the databases are screened to identify nucleic acids or polypeptides having at least 97%, at least 95%, at least 90%, at least 85%, at least 80%, at least 70%, at least 60%, or at least 50%, at least 40% identity or similarity to a nucleic acid or polypeptide involved in proliferation or an antisense nucleic acid involved in proliferation. For example, the database may be screened to identify nucleic acids homologous to one of SEQ ID Nos. 1-81, 405-485, 82-88, 90-242 or polypeptides homologous to SEQ ID NOs. 243-357, 359-398. In some embodiments, the database may be screened to identify homologous nucleic acids or polypeptides from organisms other than *E. coli*, including organisms such as *Staphylococcus aureus*, *Pseudomonas*

*aeruginosa*, *Enterobacter cloacae*, *Helicobacter pylori*, *Neisseria gonorrhoeae*,  
*Enterococcus faecalis*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Salmonella*  
*typhimurium*, *Saccharomyces cerevisiae*, *Candida albicans*, *Cryptococcus neoformans*,  
*Aspergillus fumigatus*, *Klebsiella pneumoniae*, *Salmonella typhi*, *Salmonella paratyphi*,  
5 *Salmonella choleraesuis*, *Staphylococcus epidermidis*, *Mycobacterium tuberculosis*,  
*Mycobacterium leprae*, *Treponema pallidum*, *Bacillus anthracis*, *Yersinia pestis*,  
*Clostridium botulinum*, *Campylobacter jejuni*, *Chlamydia trachomatis*, *Chlamydia*  
*pneumoniae* or any species falling within the genera of any of the above species.

10 In another embodiment, gene expression arrays and microarrays can be  
employed. Gene expression arrays are high density arrays of DNA samples deposited at  
specific locations on a glass chip, nylon membrane, or the like. Such arrays can be used  
by researchers to quantify relative gene expression under different conditions. Gene  
expression arrays are used by researchers to help identify optimal drug targets, profile  
new compounds, and determine disease pathways. An example of this technology is  
15 found in U.S. Patent No. 5807522, which is hereby incorporated by reference.

It is possible to study the expression of all genes in the genome of a particular  
microbial organism using a single array. For example, the arrays from Genosys consist  
of 12 x 24 cm nylon filters containing PCR products corresponding to 4290 ORFs from  
*E. coli*. 10 ngs of each are spotted every 1.5 mm on the filter. Single stranded labeled  
20 cDNAs are prepared for hybridization to the array (no second strand synthesis or  
amplification step is done) and placed in contact with the filter. Thus the labeled  
cDNAs are of "antisense" orientation. Quantitative analysis is done by  
phosphorimager.

Hybridization of cDNA made from a sample of total cell mRNA to such an  
25 array followed by detection of binding by one or more of various techniques known to  
those in the art results in a signal at each location on the array to which cDNA  
hybridized. The intensity of the hybridization signal obtained at each location in the  
array thus reflects the amount of mRNA for that specific gene that was present in the  
sample. Comparing the results obtained for mRNA isolated from cells grown under  
30 different conditions thus allows for a comparison of the relative amount of expression  
of each individual gene during growth under the different conditions.

Gene expression arrays may be used to analyze the total mRNA expression pattern at various time points after induction of an antisense nucleic acid against a proliferation-required gene. Analysis of the expression pattern indicated by hybridization to the array provides information on whether or not the target gene of the antisense nucleic acid is being affected by antisense induction, how quickly the antisense is affecting the target gene, and for later timepoints, what other genes are affected by antisense expression. For example, if the antisense is directed against a gene for ribosomal protein L7/L12 in the 50S subunit, its targeted mRNA may disappear first and then other mRNAs may be observed to increase, decrease or stay the same. Similarly, if the antisense is directed against a different 50S subunit ribosomal protein mRNA (e.g. L25), that mRNA may disappear first followed by changes in mRNA expression that are similar to those seen with the L7/L12 antisense expression. Thus, the mRNA expression pattern observed with an antisense nucleic acid against a proliferation required gene may identify other proliferation-required nucleic acids in the same pathway as the target of the antisense nucleic acid. In addition, the mRNA expression patterns observed with candidate drug compounds may be compared to those observed with antisense nucleic acids against a proliferation-required nucleic acid. If the mRNA expression pattern observed with the candidate drug compound is similar to that observed with the antisense nucleic acid, the drug compound may be a promising therapeutic candidate. Thus, the assay would be useful in assisting in the selection of candidate drug compounds for use in screening methods such as those described below.

In cases where the source of nucleic acid deposited on the array and the source of the nucleic acid being hybridized to the array are from two different organisms, gene expression arrays can identify homologous genes in the two organisms.

The present invention also contemplates additional methods for screening other microorganisms for proliferation-required genes. In this embodiment, the conserved portions of sequences identified as proliferation-required can be used to generate degenerate primers for use in the polymerase chain reaction (PCR). The PCR technique is well known in the art. The successful production of a PCR product using degenerate probes generated from the sequences identified herein would indicate the presence of a homologous gene sequence in the species being screened. This homologous gene is then isolated, expressed, and used as a target for candidate antibiotic compounds. In another

aspect of this embodiment, the homologous gene is expressed in an autologous organism or in a heterologous organism in such a way as to alter the level or activity of a homologous gene required for proliferation in the autologous or heterologous organism. In still another aspect of this embodiment, the homologous gene or portion is expressed in an antisense orientation in such a way as to alter the level or activity of a nucleic acid required for proliferation of an autologous or heterologous organism.

The homologous sequences to proliferation-required genes identified using the techniques described herein may be used to identify proliferation-required genes of organisms other than *E. coli*, to inhibit the proliferation of organisms other than *E. coli* by inhibiting the activity or reducing the amount of the identified homologous nucleic acid or polypeptide in the organism other than *E. coli*, or to identify compounds which inhibit the growth of organisms other than *E. coli* as described below.

In another embodiment of the present invention, *E. coli* sequences identified as required for proliferation are transferred to expression vectors capable of function within non-*E. coli* species. As would be appreciated by one of ordinary skill in the art, expression vectors must contain certain elements that are species specific. These elements can include promoter sequences, operator sequences, repressor genes, origins of replication, ribosomal binding sequences, termination sequences, and others. To use the identified exogenous sequences of the present invention, one of ordinary skill in the art would know to use standard molecular biology techniques to isolate vectors containing the sequences of interest from cultured bacterial cells, isolate and purify those sequences, and subclone those sequences into an expression vector adapted for use in the species of bacteria to be screened.

Expression vectors for a variety of other species are known in the art. For example, Cao et al. report the expression of steroid receptor fragments in *Staphylococcus aureus*. **J. Steroid Biochem Mol Biol.** 44(1):1-11 (1993). Also, Pla et al. have reported an expression vector that is functional in a number of relevant hosts including: *Salmonella typhimurium*, *Pseudomonas putida*, and *Pseudomonas aeruginosa*. **J. Bacteriol.** 172(8):4448-55 (1990). These examples demonstrate the existence of molecular biology techniques capable of constructing expression vectors for the species of bacteria of interest to the present invention.



Following the subcloning of the identified nucleic acid sequences into an expression vector functional in the microorganism of interest, the identified nucleic acid sequences are conditionally transcribed to assay for bacterial growth inhibition. Those expression vectors found to contain sequences that, when transcribed, inhibit bacterial growth are compared to the known genomic sequence of the pathogenic microorganism being screened or, if the homologous sequence from the organism being screened is not known, it may be identified and isolated by hybridization to the proliferation-required *E. coli* sequence of interest or by amplification using primers based on the proliferation-required *E. coli* sequence of interest as described above.

The antisense sequences from the second organism which are identified as described above may then be operably linked to a promoter, such as an inducible promoter, and introduced into the second organism. The techniques described herein for identifying *E. coli* genes required for proliferation may thus be employed to determine whether the identified sequences from a second organism inhibit the proliferation of the second organism.

Antisense nucleic acids required for the proliferation of organisms other than *E. coli* or the genes corresponding thereto, may also be hybridized to a microarray containing the *E. coli* ORFs to gauge the homology between the *E. coli* sequences and the proliferation-required nucleic acids from other organisms. For example, the proliferation-required nucleic acid may be from *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Helicobacter pylori*, *Neisseria gonorrhoeae*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Salmonella typhimurium*, *Saccharomyces cerevisiae*, *Candida albicans*, *Cryptococcus neoformans*, *Aspergillus fumigatus*, *Klebsiella pneumoniae*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella choleraesuis*, *Staphylococcus epidermidis*, *Mycobacterium tuberculosis*, *Mycobacterium leprae*, *Treponema pallidum*, *Bacillus anthracis*, *Yersinia pestis*, *Clostridium botulinum*, *Campylobacter jejuni* or *Chlamydia trachomatis*, *Chlamydia pneumoniae* or any species falling within the genera of any of the above species. The proliferation-required nucleic acids from an organism other than *E. coli* may be hybridized to the array under a variety of conditions which permit hybridization to occur when the probe has different levels of homology to the sequence on the

microarray. This would provide an indication of homology across the organisms as well as clues to other possible essential genes in these organisms.

In still another embodiment, the exogenous nucleic acid sequences of the present invention that are identified as required for bacterial growth or proliferation can be used as antisense therapeutics for killing bacteria. The antisense sequences can be directed against the proliferation-required genes whose sequence corresponds to the exogenous nucleic acid probes identified here (i.e. the antisense nucleic acid may hybridize to the gene or a portion thereof). Alternatively, antisense therapeutics can be directed against operons in which proliferation-required genes reside (i.e. the antisense nucleic acid may hybridize to any gene in the operon in which the proliferation-required genes reside). Further, antisense therapeutics can be directed against a proliferation-required gene or portion thereof with or without adjacent noncoding sequences, an intragenic sequence (i.e. a sequence within a gene), an intergenic sequence (i.e. a sequence between genes), a sequence spanning at least a portion of two or more genes, a 5' noncoding region or a 3' noncoding region located upstream or downstream from the actual sequence that is required for bacterial proliferation or an operon containing a proliferation-required gene.

In addition to therapeutic applications, the present invention encompasses the use of nucleic acid sequences complementary to sequences required for proliferation as diagnostic tools. For example, nucleic acid probes complementary to proliferation-required sequences that are specific for particular species of microorganisms can be used as probes to identify particular microorganism species in clinical specimens. This utility provides a rapid and dependable method by which to identify the causative agent or agents of a bacterial infection. This utility would provide clinicians the ability to prescribe species specific antimicrobial compounds to treat such infections. In an extension of this utility, antibodies generated against proteins translated from mRNA transcribed from proliferation-required sequences can also be used to screen for specific microorganisms that produce such proteins in a species-specific manner.

The following examples teach the genes of the present invention and a subset of uses for the *E. coli* genes identified as required for proliferation. These examples are illustrative only and are not intended to limit the scope of the present invention.

#### EXAMPLES

The following examples are directed to the identification and exploitation of *E. coli* genes required for proliferation. Methods of gene identification are discussed as well as a variety of methods to utilize the identified sequences.

## 5      **Genes Identified as Required for Proliferation of *E. coli***

Exogenous nucleic acid sequences were cloned into an inducible expression vector and assayed for growth inhibition activity. Example 1 describes the examination of a library of exogenous nucleic acid sequences cloned into IPTG-inducible expression vectors. Upon activation or induction, the expression vectors produced an RNA molecule  
10      corresponding to the subcloned exogenous nucleic acid sequences. The RNA product was in an antisense orientation with respect to the *E. coli* genes from which it was originally derived. This antisense RNA then interacted with sense mRNA produced from various *E. coli* genes and interfered with or inhibited the translation of the sense messenger RNA (mRNA) thus preventing protein production from these sense mRNA molecules. In cases  
15      where the sense mRNA encoded a protein required for the proliferation, bacterial cells containing an activated expression vector failed to grow or grew at a substantially reduced rate.

### **EXAMPLE 1**

#### Inhibition of Bacterial Proliferation after IPTG induction

20      To study the effects of transcriptional induction in liquid medium, growth curves were carried out by back diluting cultures 1:200 into fresh media with or without 1 mM IPTG and measuring the OD<sub>450</sub> every 30 minutes (min). To study the effects of transcriptional induction on solid medium, 10<sup>2</sup>, 10<sup>3</sup>, 10<sup>4</sup>, 10<sup>5</sup>, 10<sup>6</sup>, 10<sup>7</sup> and 10<sup>8</sup> fold dilutions of overnight cultures were prepared. Aliquots of from 0.5 to 3 µl of these  
25      dilutions were spotted on selective agar plates with or without 1 mM IPTG. After overnight incubation, the plates were compared to assess the sensitivity of the clones to IPTG.

Of the numerous clones tested, some clones were identified as a containing sequence that inhibited *E. coli* growth after IPTG induction. Accordingly, the gene to  
30      which the inserted nucleic acid sequence corresponds, or a gene within the operon containing the inserted nucleic acid, may be required for proliferation in *E. coli*.

## Characterization of Isolated Clones Negatively Affecting *E. coli* Proliferation

Following the identification of those expression vectors that, upon expression, negatively impacted *E. coli* growth or proliferation, the inserts or nucleic acid fragments contained in those expression vectors were isolated for subsequent characterization.

5 Expression vectors of interest were subjected to nucleic acid sequence determination.

### EXAMPLE 2

#### Nucleic Acid Sequence Determination of Identified Clones Expressing Nucleic Acid Fragments with Detrimental Effects of *E. coli* Proliferation

10 The nucleotide sequences for the exogenous identified sequences were determined using plasmid DNA isolated using QIAPREP (Qiagen, Valencia, CA) and methods supplied by the manufacturer. The primers used for sequencing the inserts were 5' - TGTTTATCAGACCGCTT - 3' (SEQ ID NO: 403) and 5' - ACAATTTACACAGCCTC - 3' (SEQ ID NO: 404). These sequences flank the polylinker in pLEX5BA. Sequence identification numbers (SEQ ID NOs) for the  
15 identified inserts are listed in Table I and discussed below.

### EXAMPLE 3

#### Comparison Of Isolated Sequences to Known Sequences

The nucleic acid sequences of the subcloned fragments obtained from the expression vectors discussed above were compared to known *E. coli* sequences in  
20 GenBank using BLAST version 1.4 or version 2.0.6 using the following default parameters: Filtering off, cost to open a gap=5, cost to extend a gap=2, penalty for a mismatch in the blast portion of run=-3, reward for a match in the blast portion of run=1, expectation value (e)=10.0, word size=11, number of one-line descriptions=100, number of alignments to show (B)=100. BLAST is described in Altschul, J Mol Biol. 215:403-10  
25 (1990), the disclosure of which is incorporated herein by reference in its entirety. Expression vectors were found to contain nucleic acid sequences in both the sense and antisense orientations. The presence of known genes, open reading frames, and ribosome binding sites was determined by comparison to public databases holding genetic information and various computer programs such as the Genetics Computer Group  
30 programs FRAMES and CODONPREFERENCE. Clones were designated as "antisense" if the cloned fragment was oriented to the promoter such that the RNA transcript produced was complementary to the expressed mRNA from a chromosomal locus. Clones were

designated as "sense" if they coded for an RNA fragment that was identical to a portion of a wild type mRNA from a chromosomal locus.

The sequences described in Examples 1-2 that inhibited bacterial proliferation and contained gene fragments in an antisense orientation are listed in Table I. This table lists each identified sequence by: a sequence identification number; a Molecule Number; a gene to which the identified sequence corresponds, listed according to the National Center for Biotechnology Information (NCBI), Blattner (Science 277:1453-1474(1997); also contains the *E. coli* K-12 genome sequence), or Rudd (Micro. and Mol. Rev. 62:985-1019 (1998)), (both papers are hereby incorporated by reference) nomenclatures. The CONTIG numbers for each identified sequence is shown, as well as the location of the first and last base pairs located on the *E. coli* chromosome. A Molecule Number with a "\*\*\*" indicates a clone corresponding to an intergenic sequence.

The sequences of the nucleic acid inserts of SEQ ID NOs: 1-81 from U.S. Provisional Patent Application No. 60/117,405 which inhibited proliferation were further analyzed. The reanalyzed sequences corresponding to SEQ ID NOs. 1-81 of U.S. Provisional Patent Application No. 60/117,405 have SEQ ID NOs. 405-485 in the present application.

SEQ ID NOs: 82-242 in U.S. Provisional Patent Application No. 60/117,405 are identical to SEQ ID NOs: 82-242 of the present application with the following exceptions. SEQ ID NO: 148 in the present application is the complementary strand of SEQ ID NO: 148 in U.S. Provisional Patent Application No. 60/117,405. Accordingly, the protein of SEQ ID NO: 308 which is encoded by SEQ ID NO: 148 has also been revised. SEQ ID NO: 163 in the present application is the complementary strand of SEQ ID NO: 163 in U.S. Provisional Patent Application No. 60/117,405. Accordingly, the protein of SEQ ID NO: 323 which is encoded by SEQ ID NO: 163 has also been revised.

The target gene of SEQ ID NOs. 18 and 19 of U.S. Provisional Patent Application No. 60/117,405 (SEQ ID NOs. 18, 19, 422, 423 of the present application) has been revised from *dicF* to *ftsZ* to reflect the fact that these SEQ ID NOs. include natural antisense molecules which inhibit *ftsZ* expression.

The gene products of the nucleic acids of SEQ ID NOs. 198 and 239-242 in U.S. Provisional Patent Application No. 60/117,405 and in the present application (SEQ ID

NOs. 358 and 399-402 of the present application) have been revised to reflect the fact that these nucleic acids encode nontranslated tRNAs and rRNAs. Tables I and II have been revised accordingly. The SEQ ID NOs. in Table II were also revised to reflect the fact that SEQ ID NOs: 89 and 402 were identical in U.S. Provisional Patent Application No. 60/117,405.

5

**TABLE I**  
**Identified Clones with Corresponding Genes and Operons**

SEQ ID NO.	Molecule No.	Gene (NCBI)	Gene (Blattner)	Gene (Rudd)	CONTIG
1, 405	EcXA001	<i>yhhQ</i>	<i>b3471</i>	<i>yhhQ</i>	AE000423
2, 406	EcXA002	<i>lepB</i>	<i>lepB</i>	<i>lepB</i>	AE000343
3, 407	EcXA003	<i>f586</i>	<i>b0955</i>	<i>ycbZ</i>	AE000197
4, 408	EcXA004	<i>rpsG, rpsL</i>	<i>b3341</i>	<i>rpsG, rpsL</i>	AE000410
5, 409	EcXA005a	<i>rplL, rplJ</i>	<i>b3986</i>	<i>rplL, rplJ</i>	AE000472
6, 410	EcXA005b	<i>rplL</i>	<i>rplL</i>	<i>rplL</i>	AE000472
7, 411	EcXA005c	<i>rplL, rplJ</i>	<i>rplL, rplJ</i>	<i>rplL, rplJ</i>	AE000472
8, 412	EcXA005d	<i>rplL, rplJ</i>	<i>rplL, rplJ</i>	<i>rplL, rplJ</i>	AE000472
9, 413	EcXA005e	<i>rplL</i>	<i>rplL</i>	<i>rplL</i>	AE000472
10, 414	EcXA005f	<i>rplL</i>	<i>rplL</i>	<i>rplL</i>	AE000472
11, 415	EcXA005g	<i>rplL</i>	<i>rplL</i>	<i>rplL</i>	AE000472
12, 416	EcXA006	<i>pta</i>	<i>b2297</i>	<i>pta</i>	AE000319
13, 417	EcXA007	<i>yicP</i>	<i>b3666</i>	<i>yicP</i>	AE000444
14, 418	EcXA008a	<i>yhaU</i>	<i>b3127</i>	<i>yhaU</i>	AE000394
15, 419	EcXA008b	<i>yhaU</i>	<i>yhaU</i>	<i>yhaU</i>	AE000394
16, 420	EcXA008c	<i>yhaU</i>	<i>yhaU</i>	<i>yhaU</i>	AE000394
17, 421	EcXA009	<i>ydeY</i>	<i>ydeY</i>	<i>ydeY</i>	AE000249
18, 422	EcXA010a (natural as)	<i>dicF</i>	<i>b1575</i>	<i>dicF</i>	AE000253
19, 423	EcXA010b	<i>dicF</i>	<i>dicF</i>	<i>dicF</i>	AE000253
20, 424	EcXA011	<i>fdnG</i>	<i>b1474</i>	<i>fdnG</i>	AE000244
21, 425	EcXA012a	<i>fusA</i>	<i>b3340</i>	<i>fusA</i>	AE000410
22, 426	EcXA012b	<i>fusA</i>	<i>fusA</i>	<i>fusA</i>	AE000410
23, 427	EcXA012c	<i>fusA</i>	<i>fusA</i>	<i>fusA</i>	AE000410
24, 428	EcXA013a	<i>o86</i>	<i>b2562</i>	<i>yfhL</i>	AE000342
25, 429	EcXA013b	<i>o86</i>	<i>b2562</i>	<i>yfhL</i>	AE000342
26, 430	EcXA013c	<i>o86</i>	<i>b2562</i>	<i>yfhL</i>	AE000342
27, 431	EcXA014	<i>visC</i>	<i>b2906</i>	<i>visC</i>	AE000374
28, 432	EcXA015	<i>yfdI</i>	<i>yfdI</i>	<i>yfdI</i>	AE000323
29, 433	EcXA016	<i>yeaQ</i>	<i>yeaQ</i>	<i>yeaQ</i>	AE000274
		<i>yoaG</i>	<i>yoaG</i>	<i>yoaG</i>	

SEQ ID NO.	Molecule No.	Gene (NCBI)	Gene (Blattner)	Gene (Rudd)	CONTIG
30, 434	EcXA017a	<i>yggE</i>	<i>b2922</i>	<i>yggE</i>	AE000375
31, 435	EcXA017b	<i>yggE</i>	<i>yggE</i>	<i>yggE</i>	AE000375
32, 436	EcXA018a	<i>o464</i>	<i>b2074</i>	<i>yegM</i>	AE000297
33, 437	EcXA018b	<i>o464</i>	<i>b2074</i>	<i>yegM</i>	AE000297
34, 438	EcXA019a	<i>yehA</i>	<i>yehA</i>	<i>yehA</i>	AE000300
					AE000299
35, 439	EcXA019b	<i>o172, yehA</i>	<i>o172, yehA</i>	<i>o172, yehA</i>	AE000299
36, 440	EcXA020	<i>o384, f82</i>	<i>b1794, b1795</i>	<i>ycaP, ycaQ</i>	AE000274
37, 441	EcXA021a	<i>f112</i>	<i>b0218</i>	<i>yafU</i>	AE000130
38, 442	EcXA021b	<i>f112</i>	<i>b0218</i>	<i>yafU</i>	AE000130
39, 443	EcXA022	<i>o740</i>	<i>b1629</i>	<i>ydgN</i>	AE000258
40, 444	EcXA023a	<i>f176, f382</i>	<i>b1504, b1505</i>	<i>ydeS, ydeT</i>	AE000247
41, 445	EcXA023b	<i>f176, f382</i>	<i>b1504, b1505</i>	<i>ydeS, ydeT</i>	AE000247
42, 446	EcXA024	<i>ygjM, ygjN</i>	<i>b3082</i>	<i>ygjM, ygjN</i>	AE000390
43, 447	EcXA025	<i>O2383</i>	<i>b1878</i>	<i>yecJ</i>	AE000289
44, 448	EcXA026	<i>o61</i>	Unpre- dicted	Unpre- dicted	AE000138
45, 449	EcXA027a	<i>yohH</i>	<i>yohH</i>	<i>yohH</i>	AE000303
46, 450	EcXA027b	<i>yohH</i>	<i>yohH</i>	<i>yohH</i>	AE000303
47, 451	EcXA027c	<i>yohH</i>	<i>yohH</i>	<i>yohH</i>	AE000303
		<i>yohI</i>	<i>yohI</i>	<i>yohI</i>	
48, 452	EcXA027d	<i>yohH</i>	<i>yohH</i>	<i>yohH</i>	AE000303
49, 453	EcXA028	<i>f296</i>	<i>b2305</i>	<i>yfcI</i>	AE000319
50, 454	EcXA029	<i>yjjK</i>	<i>b4391</i>	<i>yjjK</i>	AE000509
51, 455	EcXA030	<i>yi5A</i>	<i>b3557</i>	<i>yi5A</i>	AE000433
52, 456	EcXA031	<i>rplE</i>	<i>B3308</i>	<i>rplE</i>	AE000408
53, 457	EcXA032a	<i>ybgD</i>	<i>ybgD</i>	<i>ybgD</i>	AE000175
54, 458	EcXA032b **	<i>ybgD</i>	<i>ybgD</i>	<i>ybgD</i>	AE000175
		<i>gltA</i>	<i>gltA</i>	<i>gltA</i>	
55, 459	EcXA033a	<i>f477 (as)</i>	<i>b3052</i>	<i>waaE</i>	AE000387
					AE000386
56, 460	EcXA033b	<i>f477</i>	<i>b3052</i>	<i>waaE</i>	AE000387
57, 461	EcXA034a	<i>cspA</i>	<i>b3556</i>	<i>cspA</i>	AE000433
58, 462	EcXA034b	<i>cspA</i>	<i>b3556</i>	<i>cspA</i>	AE000433
59, 463	EcXA035	<i>yhjU</i>	<i>yhjU</i>	<i>yhjU</i>	AE000431
60, 464	EcXA036	<i>yqjF</i>	<i>b3101</i>	<i>yqjF</i>	AE000392
		<i>o99</i>	<i>b3100,</i>	<i>yqjK</i>	
61, 465	EcXA037	<i>ydeH</i>	<i>b1535</i>	<i>ydeH</i>	AE000251

SEQ ID NO.	Molecule No.	Gene (NCBI)	Gene (Blattner)	Gene (Rudd)	CONTIG
62, 466	EcXA038	<i>sieB</i>	<i>b1353</i>	<i>sieB</i>	AE000233
63, 467	EcXA039	<i>ybbD</i>		<i>ybbD</i>	AE000156
64, 468	EcXA040	<i>InsB_6</i>	<i>b3445</i>	<i>insB_6</i>	AE000420
65, 469	EcXA041	<i>f234</i>	<i>b1138</i>	<i>ymfE</i>	AE000214
66, 470	EcXA042a	<i>rplY</i>	<i>rplY</i>	<i>rplY</i>	AE000308
67, 471	EcXA042b	<i>rplY</i>	<i>rplY</i>	<i>rplY</i>	AE000308
68, 472	EcXA043	<i>ybgB</i>	<i>ybgB</i>	<i>ybgB</i>	AE000176
		<i>cydA</i>	<i>cydA</i>	<i>cydA</i>	
69, 473	EcXA044	<i>purB</i>	<i>b1131</i>	<i>purB</i>	AE000213
70, 474	EcXA045*	<i>csrA</i>	<i>csrA</i>	<i>csrA</i>	AE000353
	*				
		<i>serI</i>	<i>serI</i>	<i>serI</i>	
71, 475	EcXA046*	<i>fimE, fimA</i>	<i>b4313</i>	<i>fimE, fimA</i>	AE000502
	*				
72, 476	EcXA047*	<i>f96, cspB</i>	<i>f96, cspB</i>	<i>cspB, ydtS</i>	AE000252
	*				
73, 477	EcXA048	<i>yefE</i>	<i>yefE</i>	<i>yefE</i>	AE000294
74, 478	EcXA049	<i>yaiC</i>	<i>b0385</i>	<i>yaiC</i>	AE000145
75, 479	EcXA050	<i>o467, o222</i>	<i>yaiU, yaiI</i>	<i>yaiU, yaiI</i>	AE000144
76, 480	EcXA051a	<i>rplB, rplW</i>	<i>rplB, rplW</i>	<i>rplB, rplW</i>	AE000408
77, 481	EcXA051b	<i>rplW</i>	<i>rplW</i>	<i>rplW</i>	AE000408
78, 482	EcXA052	<i>infC</i>	<i>infC</i>	<i>infC</i>	AE000267
					AE000266
79, 483	EcXA053	<i>gor</i>	<i>gor</i>	<i>gor</i>	AE000426
80, 484	EcXA054	<i>rplF</i>	<i>rplF</i>	<i>rplF</i>	AE000408
81, 485	EcXA055	<i>rrlG</i>	<i>rrlG</i>	<i>rrlG</i>	AE000345

#### EXAMPLE 4

##### Identification of Genes and their Corresponding Operons Affected by Antisense Inhibition

The sequencing of the entire E. coli genome is described in Blattner et al., Science 277:1453-1474(1997) the entirety of which is hereby incorporated by reference and the sequence of the genome is listed in GenBank Accession No.U00096, the disclosure of which is incorporated herein by reference in its entirety. The operons to which the proliferation-inhibiting nucleic acids correspond were identified using RegulonDB and information in the literature. The coordinates of the boundaries of these operons on the E. coli genome are listed in Table III. Table II lists the molecule numbers of the inserts containing the growth inhibiting nucleic acid fragments, the genes in the operons



corresponding to the inserts, the SEQ ID NOs of the genes containing the inserts, the SEQ ID NOs of the proteins encoded by the genes, the start and stop points of the genes on the E. coli genome, the orientation of the genes on the genome, whether the operons are predicted or documented, and the predicted functions of the genes. The identified operons, their putative functions, and whether or not the genes are presently thought to be required for proliferation are discussed below.

Functions for the identified genes were determined by using either Blattner functional class designations or by comparing identified sequence with known sequences in various databases. A variety of biological functions were noted for the genes to which the clones of the present invention correspond. The functions for the genes of interest appear in Table II.

The proteins that are listed in Table II are involved in a wide range of biological functions.

TABLE II  
All Operon Data with Whole Chromosome Coordinates

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
82	243	EcXA001	<i>yhhQ</i>	3606848	3607513	(P)	Hypothetical ORF, unclassified, unknown	Hypothetical outer membrane protein
83	244		<i>dcrB</i>	3607532	3608143		Hypothetical ORF, unclassified, unknown	Resistance to phage C1; periplasmic protein perhaps anchored to inner membrane
84	245	EcXA002	<i>lepB</i>	2702355	2703329	(P)	Transport and binding proteins	Secretion
85	246	EcXA003	<i>ycbZ</i>	1015762	1017522	(P)	Unknown	Protease
86	247	EcXA004	<i>tufA</i>	3467782	3468966	(D)	Translation, post-translational modification	Translation (Elongation factor Tu)
87	248		<i>fusA</i>	3469037	3471151		Translation, post-translational modification	Translation (elongation factor efg)
88	249		<i>rpsG</i>	3471179	3471718		Translation, post-translational modification	Translation

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
89	402	EcXA055	<i>rrsG</i>	2727636	2729178		Translation, post-translational modification	Translation (rRNA)
90	250		<i>rpsL</i>	3471815	3471815		Translation, post-translational modification	Translation
91	251	EcXA005a-g	<i>rplJ</i>	4177574	4178071	(D)	Translation, post-translational modification	Translation
92	252		<i>rplL</i>	4178138	4178503		Translation, post-translational modification	Translation
93	253	EcXA006	<i>pta</i>	2412767	2414911	(P)	Carbon compound catabolism	Carbon compound catabolism
94	254	EcXA007	<i>yicP</i>	3841591	3843357	(P)	Hypothetical ORF, unclassified, unknown	Probable adenine deaminase
95	255	EcXA008a-c	<i>yhaD</i>	3268266	3269492	(P)	Hypothetical ORF, unclassified, unknown	
96	256		<i>yhaE</i>	3269508	3270407		Putative enzymes	
97	257		<i>yhaF</i>	3270428	3271198		Hypothetical ORF, unclassified, unknown	

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
98	258		<i>yhaU</i>	3271214	3272548		Carbon compound catabolism	Probable integral membrane protein Phthalate permease family
99	259	EcXA009	<i>ydeX</i>	1599514	1601049	(P)	Putative transport proteins	
100	260		<i>ydeY</i>	1601043	1602071		Putative transport proteins	Putative ABC transporter
101	261		<i>ydeZ</i>	1602071	1603063		Hypothetical ORF, unclassified, unknown	
102	262		<i>yneA</i>	1603075	1604097		Hypothetical ORF, unclassified, unknown	
103	263		<i>yneB</i>	1604124	1604999		Hypothetical ORF, unclassified, unknown	
104	264		<i>yneC</i>	1605023	1605313		Hypothetical ORF, unclassified, unknown	
105	265	EcXA010a-b	<i>fisZ</i>	105305	106456	(P)	Cell processes (incl. Adaptation, protection)	Regulator of cell division
106	266	EcXA011	<i>fdnG</i>	1545425	1548472	(D)	Energy metabolism	Anaerobic respiration (formate dehydro-

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
								genase)
107	267		<i>fhnH</i>	1548485	1549369		Energy metabolism	
108	268		<i>fhnI</i>	1549362	1550015		Energy metabolism	
		EcXA 012a-c	Same operon as EcXA004					
109	269	EcXA013a-c	<i>yhlL</i>	2697683	2697943	(P)	Hypothetical ORF, unclassified, unknown	No homologues, no motifs
110	270	EcXA014	<i>visC</i>	3049135	3050337	(P)	Hypothetical ORF, unclassified, unknown	Ubiquinone synthesis
111	271		<i>ubiH</i>	3050360	3051538		Biosynthesis of cofactors, prosthetic groups and carriers	
112	272		<i>pepP</i>	3051535	3052860		Translation, post-translational modification	
113	273		<i>yglB</i>	3052886	3053470		Hypothetical ORF, unclassified, unknown	
114	274	EcXA015	<i>yglG</i>	2465875	2466237	(P)	Hypothetical ORF, unclassified, unknown	

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
115	275		<i>yflH</i>	2466234	2467154		Cell structure	
116	276		<i>yflI</i>	2467151	2468482		Hypothetical ORF, unclassified, unknown	Putative membrane protein
117	277	EcXA016	<i>yeaQ</i>	1877031	1877279	(P)	Hypothetical ORF, unclassified, unknown	Homologue to transglycosylase associated protein
118	278		<i>yoaG</i>	1877427	1877609	(P)	Hypothetical ORF, unclassified, unknown	No homologues
119	279		<i>yeaR</i>	1877613	1877972		Hypothetical ORF, unclassified, unknown	
120	280	EcXA017a-b	<i>yggE</i>	3065360	3066100	(P)	Structural proteins	Homologues in multiple bacteria, no motifs
121	281	EcXA018a-b	<i>yegM</i>	2151891	2153285	(P)	Putative transport proteins	Transport (multiple transferable resistance)
122	282		<i>yegN</i>	2153285	2156407		Hypothetical ORF, unclassified, unknown	
123	283		<i>yegO</i>	2156408	2159485		Hypothetical ORF, unclassified, unknown	

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
124	284		<i>yegB</i>	2159486	2160901		Putative transport proteins	
125	285	EcXA019a-b	<i>yehA</i>	2185400	2186434	(P)	Cell structure	Weak homology to pilin precursor from <i>H. Inf.</i>
126	286		<i>yehB</i>	2186450	2188930		Hypothetical ORF, unclassified, unknown	
127	287		<i>yehC</i>	2188946	2189665		Putative chaperones	
128	288		<i>yehD</i>	2189700	2190242		Cell structure	
		EcXA020	Same operon as EcXA016 (one of the two)					
129	289	EcXA021a-b	<i>yafU</i>	238746	239084	(P)	Hypothetical ORF, unclassified, unknown	Homologues in <i>H. Inf.</i> and <i>S. Pombe.</i> , no motifs, transmembrane region present
130	290	EcXA022	<i>ydgL</i>	1703791	1704372	(P)	Hypothetical ORF, unclassified, unknown	
131	291		<i>ydgM</i>	1704372	1704950		Hypothetical ORF, unclassified, unknown	

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
132	292		<i>ydgN</i>	1704943	1707165		Hypothetical ORF, unclassified, unknown	
133	293		<i>ydgO</i>	1707166	1708224		Hypothetical ORF, unclassified, unknown	
134	294		<i>ydgP</i>	1708228	1708848		Hypothetical ORF, unclassified, unknown	
135	295		<i>ydgQ</i>	1708852	1709547		Hypothetical ORF, unclassified, unknown	
136	296		<i>nth</i>	1709547	1710182		Transcription, RNA processing and degradation	
137	297	EcXA023a-b	<i>ydeR</i>	1585817	1586320	(P)	Hypothetical ORF, unclassified, unknown	
138	298		<i>ydeS</i>	1586333	1586863		Hypothetical ORF, unclassified, unknown	fimf-like
139	299		<i>ydeT</i>	1586877	1588025		Structural proteins	fimd-like
140	300	EcXA024	<i>yglM</i>	3231369	3231785	(P)	Hypothetical ORF, unclassified, unknown	Weak homology to long chain fatty acid coa ligase in



GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
								Archaeoglobus
141	301		<i>ygiN</i>	3231782	3232096		Hypothetical ORF, unclassified, unknown	Homologues in various bacteria
142	302	EcXA025	<i>yecJ</i>	2042885	2050036	(P)	Hypothetical ORF, unclassified, unknown	Strong similarity to numerous attaching and effacing proteins and invasins
143	303	EcXA026	<i>yajA</i>	331001	331184	unpredicted		nifm like
144	304	EcXA027a-d	<i>yohG</i>	2225343	2226539	(P)	Putative transport proteins	
145	305		<i>yohH</i>	2226569	2226859		Hypothetical ORF, unclassified, unknown	Xylose binding protein-like
146	306		<i>yohl</i>	2227458	2228405	(P)	Putative regulatory protein	
147	307	EcXA028	<i>yefl</i>	2420669	2421559	(P)	Hypothetical ORF, unclassified, unknown	Similar to <i>S. Typhi</i> histidine transport gene
148	308	EcXA029	<i>yjiK</i>	4626424	4628091	(P)	Hypothetical ORF, unclassified, unknown	Similar to ABC transporter

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
149	309	EcXA030	<i>yi5A</i>	3718309	3718830	(P)	Hypothetical ORF, unclassified, unknown	IS150 orf A
150	310		<i>yi5B</i>	3718827	3719678		Phage, transposon, or plasmid	
151	311	EcXA031	<i>rpmJ</i>	3440255	3440371	(D)	Translation, post-translational modification	
152	312		<i>prlA</i>	3440403	3441734		Putative transport proteins	
153	313		<i>rplO</i>	3441742	3442176		Translation, post-translational modification	
154	314		<i>rpmD</i>	3442180	3442359		Translation, post-translational modification	
155	315		<i>rpsE</i>	3442363	3442866		Translation, post-translational modification	
156	316		<i>rplR</i>	3442881	3443234		Translation, post-translational modification	
157	317		<i>rplF</i>	3443244	3443777		Translation, post-translational modification	Translation

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
158	318		<i>rpsH</i>	3443790	3444182		Translation, post-translational modification	
159	319		<i>rpsN</i>	3444216	3444521		Translation, post-translational modification	
160	320		<i>rplE</i>	3444536	3445075		Translation, post-translational modification	Translation
161	321		<i>rplX</i>	3445090	3445404		Translation, post-translational modification	
162	322		<i>rplN</i>	3445415	3445786		Translation, post-translational modification	
163	323	EcXA032a-b	<i>ybgD</i>	751452	752018	(P)	Cell processes (incl. Adaptation, protection)	Hypothetical fimbrial protein
164	324		<i>gluA</i>	752408	753691	(D)	Energy metabolism	Glutamine biosynthesis
165	325	EcXA033a-b	<i>waaE</i>	3192961	3194394	(P)	Putative enzymes	ADP heptose synthase/ autotrophic growth protein
166	326		<i>glnE</i>	3194442	3197282		Translation, post-translational	

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
							modification	
167	327		<i>ygiF</i>	3197305	3198606		Hypothetical ORF, unclassified, unknown	
168	328	EcXA034a-b	<i>cspA</i>	3717678	3717890	(P)	Cell processes (incl. Adaptation, protection)	RNA chaperonin
169	329	EcXA035	<i>yhjS</i>	3694087	3695658	(P)	Translation, post-translational modification	
170	330		<i>yhjT</i>	3695658	3695846		Hypothetical ORF, unclassified, unknown	
171	331		<i>yhjU</i>	3695843	3697522		Hypothetical ORF, unclassified, unknown	Regions similar to dehydro-genases, nucleases etc.
172	332	EcXA036	<i>yqiC</i>	3246594	3246977	(P)	Hypothetical ORF, unclassified, unknown	
173	333		<i>yqiD</i>	3247015	3247320		Hypothetical ORF, unclassified, unknown	
174	334		<i>yqiE</i>	3247323	3247727		Hypothetical ORF, unclassified,	

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
							unknown	
175	335		<i>yqjK</i>	3247717	3248016		Similar to mukB from <i>H. Inf.</i>	
176	336		<i>yqjF</i>	3248112	3248594	(P)	Hypothetical ORF, unclassified, unknown	Homologues in many bacteria, blocks; secretion/ ATP synthase/ftsZ
177	337	EcXA037	<i>ydeH</i>	1620984	1621874	(P)	Hypothetical ORF, unclassified, unknown	Similar to carboxy-kinase, oxidase, symporters
178	338	EcXA038	<i>sieB</i>	1416572	1417183	(P)	Phage, transposon, or plasmid	Super-infection exclusion factor B-like
179	339		<i>rajB</i> ( <i>h1354</i> )	1417192	1417368		Hypothetical ORF, unclassified, unknown	
180	340	EcXA039	<i>rhsD</i>	522485	526765	(P)	Hypothetical ORF, unclassified, unknown	
181	341		<i>ybbC</i>	526805	527173		Hypothetical ORF, unclassified, unknown	
182	342		<i>ybhH</i>	527173	527883		Hypothetical ORF, unclassified, unknown	Rhs-like element

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
							unknown	
183	343		<i>ybbD</i>	527864	528124		Hypothetical ORF, unclassified, unknown	ATP synthase, desaturase
184	344		<i>yblI</i>	528163	528354		Hypothetical ORF, unclassified, unknown	
185	345	EcXA040	<i>insB_6</i>	351114	351389	(P)	Phage, transposon, or plasmid	
186	346		<i>insA</i>	351308	3581811		Phage, transposon, or plasmid	
187	347		<i>yrhA</i>	3580669	3581085		Hypothetical ORF, unclassified, unknown	
188	348		<i>yhhZ</i>	3579494	3580672		Hypothetical ORF, unclassified, unknown	
189	349	EcXA041	<i>ymfD</i>	1196090	1196755	(P)	Hypothetical ORF, unclassified, unknown	No assigned role
190	350		<i>ymfE</i>	1196756	1197460		Hypothetical ORF, unclassified, unknown	No assigned role

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
191	351	EcXA042a-b	<i>rplY</i>	2280537	2280821	(P)	Translation, post-translational modification	Translation
192	352	EcXA043	<i>hrsA</i>	765207	767183	(P)	Translation, post-translational modification	
193	353		<i>ybgB</i>	767201	769834		Carbon compound catabolism	Unknown
194	354		<i>cydA</i>	770678	772249	(D)	Energy metabolism	Cytochrome D oxidase
195	355		<i>cydB</i>	772265	773404		Energy metabolism	
196	356	EcXA044	<i>purB</i>	1189839	1191209	(D)	Nucleotide biosynthesis and metabolism	Purine biosynthesis
197	357	EcXA045	<i>csrA</i>	2816983	2817168	(P)	Regulatory function	Carbon storage regulator (mRNA decay factor)
198	358		<i>serV</i>	2816575	2816667	Unpredicted	Translation, post-translational modification	Translation (tRNA)
199	359	EcXA046	<i>fimB</i>	4538525	4539127	(D)	Cell structure	
200	360		<i>fimE</i>	4539605	4540201		Cell structure	Fimbrac
201	361		<i>fimA</i>	4540683	4541231		Cell structure	Regulator of inversion
202	362		<i>fimI</i>	4541188	4541835		Cell structure	

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
203	363		<i>fimC</i>	4541872	4542597		Cell structure	
204	364		<i>fimD</i>	4542665	4545301		Cell structure	
205	365		<i>fimF</i>	4545311	4545841		Cell structure	
206	366		<i>fimG</i>	4545854	4546357		Cell structure	
207	367		<i>fimH</i>	4546377	4547279		Cell structure	
208	368	EcXA047	<i>ydfP</i>	1637054	1638684	(P)	Hypothetical ORF, unclassified, unknown	
209	369		<i>ydfQ</i>	1637548	1638081		Hypothetical ORF, unclassified, unknown	
210	370		<i>ydfR</i>	1638078	1638389		Hypothetical ORF, unclassified, unknown	
211	371		<i>ydfS</i>	1638394	1638684		Hypothetical ORF, unclassified, unknown	Lysis protein
212	372		<i>cspB</i>	1639363	1639578	(P)	Cell processes (incl. Adaptation, protection)	
213	373	EcXA048	<i>yi52_7</i>	2099917	2100933	(P)	Phage, transposon, or plasmid	
214	374		<i>yefJ</i>	2100938	2101411		Putative enzymes	
215	375		<i>yefI</i>	2101413	2102531		Hypothetical ORF, unclassified,	



GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
							unknown	
216	376		<i>yefH</i>	2102516	2103106		Putative enzymes	
217	377		<i>yefG</i>	2103087	2104079		Hypothetical ORF, unclassified, unknown	
218	378		<i>rfc</i>	2104082	2105248		Cell structure	
219	379		<i>yefE</i>	2105248	2106351		Hypothetical ORF, unclassified, unknown	UDP galactopyranase mutase
220	380	EcXA049	<i>yaiC</i>	402927	404042	(P)	Hypothetical ORF, unclassified, unknown	Unknown
221	381	EcXA050	<i>yaiU</i>	392239	393642	(P)	Putative enzymes	Putative auto-transporter
222	382		<i>yaiV</i>	393685	394353		Hypothetical ORF, unclassified, unknown	Hypothetical outer membrane protein
223	383	EcXA051a-b	<i>rpsQ</i>	3445951	3446205	(D)	Translation, post-translational modification	
224	384		<i>rpmC</i>	3446205	3446396		Translation, post-translational modification	



GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
233	393		<i>rpsJ</i>	3450596	3450907		Translation, post-translational modification	
234	394	EcXA052	<i>rplT</i>	1797417	1797773	(D)	Translation, post-translational modification	
235	395		<i>rpmI</i>	1797826	1798023		Translation, post-translational modification	
236	396		<i>infC</i>	1798120	1798662		Translation, post-translational modification	Translation
237	397		<i>thrS</i>	1798666	1800594		Translation, post-translational modification	
238	398	EcXA053	<i>gor</i>	3643929	3645281	(P)	Biosynthesis of cofactors, prosthetic groups and carriers	Glutathione oxidoreductase
		EcXA054	Same operon as EcXA031					
239	399	EcXA055	<i>rrlG</i>	2724301	2727204	(D)	Translation, post-translational modification	Translation (rRNA)

GeneSeq ID No.	Gene Prod. Seq ID No.	Mole. No.	Genes On Operon	Left Coordinate	Right Coordinate	Predicted (P) Or Documented (D) Operon	Blattner functional class of encoded proteins	Predicted functional class of encoded proteins
240	400		<i>rrfG</i>	2724089	2724208		Translation, post-translational modification	Translation (rRNA)
241	401		<i>gltW</i>	2727389	2727464		Translation, post-translational modification	Translation (tRNA)
242	402		<i>rrsG</i>	2727636	2729178		Translation, post-translational modification	Translation (rRNA)

Several of the expression vectors contain fragments that correspond to genes of unknown function or if the function is known, it is not known whether the gene is essential. For example, EcXA001, 003, 007, 008, 013, 015, 016, 017, 018, 019, 020, 021, 022, 023, 024, 025, 026, 027, 028, 029, 030, 032, 033, 034, 035, 036, 037, 038, 039, 040, 041, 047, 048, 049 and 050 are all exogenous nucleic acid sequences that correspond to *E. coli* proteins that have no known function or where the function has not been shown to be essential or nonessential.

The present invention reports a number of novel *E. coli* genes and operons that are required for proliferation. From the list clone sequences identified here, each was identified to be a portion of a gene in an operon required for the proliferation of *E. coli*. Cloned sequences corresponding to genes already known to be required for proliferation in *E. coli* include EcXA002, 004, 005, 010, 012, 014, 031, 02, 043, 045, 051, 052, 054, and 055. The remaining identified sequences correspond to *E. coli* genes previously undesignedated as required for proliferation in the art.

An interesting observation of the present invention is that there are also several sequence fragments that correspond to *E. coli* genes that are not thought to be required for *E. coli* proliferation. Nevertheless, under the conditions described above, the antisense expression of these gene fragments causes a reduction in cell growth. This result implies that the genes corresponding to the identified sequences are actually required for proliferation. Molecule Nos. corresponding to these genes are EcXA006, 044, 046, and 053.

Following identification of the sequences of interest, these sequences were localized into operons. Since bacterial genes are expressed in a polycistronic manner, the antisense inhibition of a single gene in an operon might effect the expression of all the other genes on the operon or the genes down stream from the single gene identified. In order to determine which of the gene products in an operon are required for proliferation, each of the genes contained within an operon may be analyzed for their effect on viability as described below.

TABLE III  
Operon Boundaries

Mole. No.	Left Coordinate	Right Coordinate
EcXA001	3606848	3608143
EcXA002	2702355	2703329
EcXA003	1015762	1017522
EcXA004	3467782	3472189
EcXA005	4177574	4178503
EcXA006	2412767	2414911
EcXA007	3841591	3843357
EcXA008	3268266	3272548
EcXA009	1599514	1605313
EcXA010	1647406	1647458
EcXA011	1545425	1550015
EcXA012	3467782	3472189
EcXA013	2697683	2697943
EcXA014	3049135	3053470
EcXA015	2465875	2468482
EcXA016	1877031	1877972
EcXA017	3065360	3066100
EcXA018	2151891	2160901
EcXA019	2185400	2190242
EcXA020	1877031	1877972
EcXA021	238746	239084
EcXA022	1703791	1710182
EcXA023	1585817	1588025
EcXA024	3231369	3232096
EcXA025	2042885	2050036
EcXA026	331001	331184
EcXA027c	2225343	2228405
EcXA028	2420669	2421559
EcXA029	4626424	4628091
EcXA030	3718309	3719678
EcXA031	3440255	3445786
EcXA032b	751452	753691
EcXA033	3192961	3198606
EcXA034	3717678	3717890
EcXA035	3694087	3697522
EcXA036	3246594	3248594
EcXA037	1620984	1621874
EcXA038	1416572	1417368
EcXA039	522485	528354
EcXA040	3580669	3580672
EcXA041	1196090	1197460

Mole. No.	Left Coordinate	Right Coordinate
EcXA042	2280537	2280821
EcXA043	765207	773404
EcXA044	1189839	1191209
EcXA045	2816575	2817168
EcXA046	4538525	4547279
EcXA047	1637054	1639578
EcXA048	2099917	2106351
EcXA049	402927	404042
EcXA050	392239	394353
EcXA051	3445951	3450907
EcXA052	1797417	1800594
EcXA053	3643929	3645281
EcXA054	3440255	3445786
EcXA055	2724301	2729178

### EXAMPLE 5

#### Identification of Individual Genes within an Operon Required for Proliferation

5 The following example illustrates a method for determining which gene in an operon is required for proliferation. The clone insert corresponding to Molecule No. EcXA004 possesses nucleic acid sequence homology to the *E. coli* genes *rspG* and *rspL*. This molecule corresponds to an operon containing two additional genes *fusA* and *tufA*. The *rpsL* gene is the first gene in the operon. To determine which gene or genes in this operon are required for proliferation, each gene is selectively inactivated using  
10 homologous recombination. Gene *rpsL* is the first gene to be inactivated.

Deletion inactivation of a chromosomal copy of a gene in *E. coli* can be accomplished by integrative gene replacement. The principle of this method (Hamilton, C. M., et al 1989. *J. Bacteriol.* **171**: 4617-4622) is to construct a mutant allele of the targeted gene, introduce that allele into the chromosome using a conditional suicide  
15 vector, and then force the removal of the native wild type allele and vector sequences. This will replace the native gene with a desired mutation(s) but leave promoters, operators, etc. intact. Essentiality of a gene is determined either by deduction from genetic analysis or by conditional expression of a wild type copy of the targeted gene (trans complementation).

20 The first step is to generate a mutant *rpsL* allele using PCR amplification. Two sets of PCR primers are chosen to produce a copy of *rpsL* with a large central deletion

to inactivate the gene. In order to eliminate polar effects, it is desirable to construct a mutant allele comprising an in-frame deletion of most or all of the coding region of the *rpsL* gene. Each set of PCR primers is chosen such that a region flanking the gene to be amplified is sufficiently long to allow recombination (typically at least 500 nucleotides on each side of the deletion). The targeted deletion or mutation will be contained within this fragment. To facilitate cloning of the PCR product, the PCR primers may also contain restriction endonuclease sites found in the cloning region of a conditional knockout vector such as pKO3 (Link, et al 1997 *J. Bacteriol.* **179** (20): 6228-6237). Suitable sites include NotI, SalI, BamHI and SmaI. The *rpsL* gene fragments are produced using standard PCR conditions including, but not limited to, those outlined in the manufacturers directions for the Hot Start Taq PCR kit (Qiagen, Inc., Valencia, CA). The PCR reactions will produce two fragments that can be fused together. Alternatively, crossover PCR can be used to generate a desired deletion in one step (Ho, S. N., et al 1989. *Gene* **77**: 51-59, Horton, R. M., et al 1989. *Gene* **77**: 61-68). The mutant allele thus produced is called a "null" allele because it cannot produce a functional gene product.

The mutant allele obtained from PCR amplification is cloned into the multiple cloning site of pKO3. Directional cloning of the *rpsL* null allele is not necessary. The pKO3 vector has a temperature-sensitive origin of replication derived from pSC101. Therefore, clones are propagated at the permissive temperature of 30°C. The vector also contains two selectable marker genes: one that confers resistance to chloramphenicol and another, the *Bacillus subtilis* *sacB* gene, that allows for counter-selection on sucrose containing growth medium. Clones that contain vector DNA with the null allele inserted are confirmed by restriction endonuclease analysis and DNA sequence analysis of isolated plasmid DNA. The plasmid containing the *rpsL* null allele insert is known as a knockout plasmid.

Once the knockout plasmid has been constructed and its sequence verified, it is transformed into a Rec<sup>+</sup> *E. coli* host cell. Transformation can be by any standard method such as electroporation. In some fraction of the transformed cells, plasmids will integrate into the *E. coli* chromosome by homologous recombination between the *rpsL* null allele in the plasmid and the *rpsL* gene in the chromosome. Transformant colonies in which such an event has occurred are readily selected by growth at the non-



permissive temperature of 43°C and in the presence of chloramphenicol. At this temperature, the plasmid will not replicate as an episome and will be lost from cells as they grow and divide. These cells are no longer resistant to chloramphenicol and will not grow when it is present. However, cells in which the knockout plasmid has integrated into the *E. coli* chromosome remain resistant to chloramphenicol and propagate.

Cells containing integrated knock-out plasmids are usually the result of a single crossover event that creates a tandem repeat of the mutant and native wild type alleles of *rpsL* separated by the vector sequences. A consequence of this is that *rpsL* will still be expressed in these cells. In order to determine if the gene is essential for growth, the wild type copy must be removed. This is accomplished by selecting for plasmid excision, a process in which homologous recombination between the two alleles results in looping out of the plasmid sequences. Cells that have undergone such an excision event and have lost plasmid sequences including *sacB* gene are selected for by addition of sucrose to the medium. The *sacB* gene product converts sucrose to a toxic molecule. Thus counter selection with sucrose ensures that plasmid sequences are no longer present in the cell. Loss of plasmid sequences is further confirmed by testing for sensitivity to chloramphenicol (loss of the chloramphenicol resistance gene). The latter test is important because occasionally a mutation in the *sacB* gene can occur resulting in a loss of *sacB* function with no effect on plasmid replication (Link, et. al., 1997 *J. Bacteriol.* **179** (20): 6228-6237). These artifact clones retain plasmid sequences and are therefore still resistant to chloramphenicol.

In the process of plasmid excision, one of the two *rpsL* alleles is lost from the chromosome along with the plasmid DNA. In general, it is equally likely that the null allele or the wild type allele will be lost. Therefore, if the *rpsL* gene is not essential, half of the clones obtained in this experiment will have the wild type allele on the chromosome and half will have the null allele. However, if the *rpsL* gene is essential, cells containing the null allele will not be obtained as a single copy of the null allele would be lethal.

To determine the essentiality of *rpsL*, a statistically significant number of the resulting clones, at least 20, are analyzed by PCR amplification of the *rpsL* gene. Since the null allele is missing a significant portion of the *rpsL* gene, its PCR product is

significantly shorter than that of the wild type gene and the two are readily distinguished by gel electrophoretic analysis. The PCR products may also be subjected to sequence determination for further confirmation by methods well known to those in the art.

5           The above experiment is generally adequate for determining the essentiality of a gene such as *rpsL*. However, it may be necessary or desirable to more directly confirm the essentiality of the gene. There are several methods by which this can be accomplished. In general, these involve three steps: 1) construction of an episome containing a wild type allele, 2) isolation of clones containing a single chromosomal  
10 copy of the mutant null allele as described above but in the presence of the episomal wild type allele, and then 3) determining if the cells survive when the expression of the episomal allele is shut off. In this case, the trans copy of wild type *rpsL* is made by PCR cloning of the entire coding region of *rpsL* and inserting it in the sense orientation downstream of an inducible promoter such as the *E. coli lac* promoter. Transcription of  
15 this allele of *rpsL* will be induced in the presence of IPTG which inactivates the *lac* repressor. Under IPTG induction *rpsL* protein will be expressed as long as the recombinant gene also possesses a ribosomal binding site, also known as a "Shine-Dalgarno Sequence". The trans copy of *rpsL* is cloned on a plasmid that is compatible with pSC101. Compatible vectors include p15A, pBR322, and the pUC plasmids,  
20 among others. Replication of the compatible plasmid will not be temperature-sensitive. The entire process of integrating the null allele of *rpsL* and subsequent plasmid excision is carried out in the presence of IPTG to ensure the expression of functional *rpsL* protein is maintained throughout. After the null *rpsL* allele is confirmed as integrated on the chromosome in place of the wild type *rpsL* allele, then IPTG is withdrawn and  
25 expression of functional *rpsL* protein shut off. If the *rpsL* gene is essential, cells will cease to proliferate under these conditions. However, if the *rpsL* gene is not essential, cells will continue to proliferate under these conditions. In this experiment, essentiality is determined by conditional expression of a wild type copy of the gene rather than inability to obtain the intended chromosomal disruption.

30           An advantage of this method over some other gene disruption techniques is that the targeted gene can be deleted or mutated without the introduction of large segments of foreign DNA. Therefore, polar effects on downstream genes are eliminated or

minimized. There are methods described to introduce inducible promoters upstream of potential essential bacterial genes. However in such cases, polarity from multiple transcription start points can be a problem. One way of preventing this is to insert a gene disruption cassette that contains strong transcriptional terminators upstream of the integrated inducible promoter (Zhang, Y, and Cronan, J. E. 1996 *J. Bacteriol.* **178** (12): 3614-3620). The described techniques will all be familiar to one of ordinary skill in the art.

Following the analysis of the *rpsL* gene, the other genes of the operon are investigated to determine if they are required for proliferation.

#### EXAMPLE 6

##### Expression of the Proteins Encoded by Genes Identified as Required for *E. coli* Proliferation

The following is provided as one exemplary method to express the proliferation-required proteins encoded by the identified sequences described above. First, the initiation and termination codons for the gene are identified. If desired, methods for improving translation or expression of the protein are well known in the art. For example, if the nucleic acid encoding the polypeptide to be expressed lacks a methionine codon to serve as the initiation site, a strong Shine-Delgarno sequence, or a stop codon, these sequences can be added. Similarly, if the identified nucleic acid sequence lacks a transcription termination signal, this sequence can be added to the construct by, for example, splicing out such a sequence from an appropriate donor sequence. In addition, the coding sequence may be operably linked to a strong promoter or an inducible promoter if desired. The identified nucleic acid sequence or portion thereof encoding the polypeptide to be expressed is obtained by PCR from the bacterial expression vector or genome using oligonucleotide primers complementary to the identified nucleic acid sequence or portion thereof and containing restriction endonuclease sequences for *NcoI* incorporated into the 5' primer and *BglII* at the 5' end of the corresponding 3'-primer, taking care to ensure that the identified nucleic acid sequence is positioned in frame with the termination signal. The purified fragment obtained from the resulting PCR reaction is digested with *NcoI* and *BglII*, purified and ligated to an expression vector.

The ligated product is transformed into DH5 $\alpha$  or some other *E. coli* strain suitable for the over expression of potential proteins. Transformation protocols are well known in the art. For example, transformation protocols are described in: **Current Protocols in**

**Molecular Biology**, Vol. 1, Unit 1.8, (Ausubel, et al., Eds.) John Wiley & Sons, Inc. (1997). Positive transformants are selected after growing the transformed cells on plates containing 50-100 µg/ml Ampicillin (Sigma, St. Louis, Missouri). In one embodiment, the expressed protein is held in the cytoplasm of the host organism. In an alternate  
5 embodiment, the expressed protein is released into the culture medium. In still another alternative, the expressed protein can be sequestered in the periplasmic space and liberated therefrom using any one of a number of cell lysis techniques known in the art. For example, the osmotic shock cell lysis method described in Chapter 16 of **Current Protocols in Molecular Biology**, Vol. 2, (Ausubel, et al., Eds.) John Wiley & Sons, Inc.  
10 (1997). Each of these procedures can be used to express a proliferation-required protein.

Expressed proteins, whether in the culture medium or liberated from the periplasmic space or the cytoplasm, are then purified or enriched from the supernatant using conventional techniques such as ammonium sulfate precipitation, standard  
15 chromatography, immunoprecipitation, immunochromatography, size exclusion chromatography, ion exchange chromatography, and HPLC. Alternatively, the secreted protein can be in a sufficiently enriched or pure state in the supernatant or growth media of the host to permit it to be used for its intended purpose without further enrichment. The purity of the protein product obtained can be assessed using techniques such as Coomassie or silver staining or using antibodies against the control protein. Coomassie and silver  
20 staining techniques are familiar to those skilled in the art.

Antibodies capable of specifically recognizing the protein of interest can be generated using synthetic peptides using methods well known in the art. See, **Antibodies: A Laboratory Manual**, (Harlow and Lane, Eds.) Cold Spring Harbor Laboratory (1988). For example, 15-mer peptides having a sequence encoded by the appropriate identified  
25 gene sequence of interest or portion thereof can be chemically synthesized. The synthetic peptides are injected into mice to generate antibodies to the polypeptide encoded by the identified nucleic acid sequence of interest or portion thereof. Alternatively, samples of the protein expressed from the expression vectors discussed above can be purified and subjected to amino acid sequencing analysis to confirm the identity of the recombinantly  
30 expressed protein and subsequently used to raise antibodies. An Example describing in detail the generation of monoclonal and polyclonal antibodies appears in Example 7.

The protein encoded by the identified nucleic acid sequence of interest or portion thereof can be purified using standard immunochromatography techniques. In such procedures, a solution containing the secreted protein, such as the culture medium or a cell extract, is applied to a column having antibodies against the secreted protein attached to the chromatography matrix. The secreted protein is allowed to bind the immunochromatography column. Thereafter, the column is washed to remove non-specifically bound proteins. The specifically bound secreted protein is then released from the column and recovered using standard techniques. These procedures are well known in the art.

In an alternative protein purification scheme, the identified nucleic acid sequence of interest or portion thereof can be incorporated into expression vectors designed for use in purification schemes employing chimeric polypeptides. In such strategies the coding sequence of the identified nucleic acid sequence of interest or portion thereof is inserted in-frame with the gene encoding the other half of the chimera. The other half of the chimera can be maltose binding protein (MBP) or a nickel binding polypeptide encoding sequence. A chromatography matrix having antibody to MBP or nickel attached thereto is then used to purify the chimeric protein. Protease cleavage sites can be engineered between the MBP gene or the nickel binding polypeptide and the identified expected gene of interest, or portion thereof. Thus, the two polypeptides of the chimera can be separated from one another by protease digestion.

One useful expression vector for generating maltose binding protein fusion proteins is pMAL (New England Biolabs), which encodes the *malE* gene. In the pMal protein fusion system, the cloned gene is inserted into a pMal vector downstream from the *malE* gene. This results in the expression of an MBP-fusion protein. The fusion protein is purified by affinity chromatography. These techniques as described are well known to those skilled in the art of molecular biology.

#### EXAMPLE 7

##### Production of an Antibody to an isolated *E. coli* Protein

Substantially pure protein or polypeptide is isolated from the transformed cells as described in Example 6. The concentration of protein in the final preparation is adjusted, for example, by concentration on a 10,000 molecular weight cut off AMICON filter device

(Millipore, Bedford, MA). to the level of a few micrograms/ml. Monoclonal or polyclonal antibody to the protein can then be prepared as follows:

### **Monoclonal Antibody Production by Hybridoma Fusion**

5 Monoclonal antibody to epitopes of any of the peptides identified and isolated as described can be prepared from murine hybridomas according to the classical method of Kohler, G. and Milstein, C., **Nature** **256**:495 (1975) or any of the well-known derivative methods thereof. Briefly, a mouse is repetitively inoculated with a few micrograms of the selected protein or peptides derived therefrom over a period of a few weeks. The mouse is then sacrificed, and the antibody producing cells of the spleen isolated. The spleen cells  
10 are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused cells destroyed by growth of the system on selective media comprising aminopterin (HAT media). The successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as described by Engvall, E.,  
15 "Enzyme immunoassay ELISA and EMIT," **Meth. Enzymol.** **70**:419 (1980), and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Davis, L. et al. **Basic Methods in Molecular Biology** Elsevier, New York. Section 21-2.  
20

### **Polyclonal Antibody Production by Immunization**

Polyclonal antiserum containing antibodies to heterogeneous epitopes of a single protein or a peptide can be prepared by immunizing suitable animals with the expressed protein or peptides derived therefrom described above, which can be unmodified or  
25 modified to enhance immunogenicity. Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. For example, small molecules tend to be less immunogenic than larger molecules and can require the use of carriers and adjuvant. Also, host animals vary in response to site of inoculations and dose, with both inadequate or excessive doses of antigen resulting in low titer antisera.  
30 Small doses (ng level) of antigen administered at multiple intradermal sites appears to be most reliable. An effective immunization protocol for rabbits can be found in Vaitukaitis, J. et al. **J. Clin. Endocrinol. Metab.** **33**:988-991 (1971).

Booster injections can be given at regular intervals, and antiserum harvested when antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against known concentrations of the antigen, begins to fall. See, for example, Ouchterlony, O. et al., Chap. 19 in: **Handbook of Experimental Immunology** D. Wier (ed) Blackwell (1973). Plateau concentration of antibody is usually in the range of 0.1 to 0.2 mg/ml of serum (about 12  $\mu$ M). Affinity of the antisera for the antigen is determined by preparing competitive binding curves, as described, for example, by Fisher, D., Chap. 42 in: **Manual of Clinical Immunology**, 2d Ed. (Rose and Friedman, Eds.) Amer. Soc. For Microbiol., Washington, D.C. (1980).

Antibody preparations prepared according to either protocol are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies can also be used in therapeutic compositions for killing bacterial cells expressing the protein.

## EXAMPLE 8

### Screening Chemical Libraries

#### A. Protein-Based Assays

Having isolated and expressed bacterial proteins shown to be required for bacterial proliferation, the present invention further contemplates the use of these expressed proteins in assays to screen libraries of compounds for potential drug candidates. The generation of chemical libraries is well known in the art. For example combinatorial chemistry can be used to generate a library of compounds to be screened in the assays described herein. A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis by combining a number of chemical "building blocks" reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining amino acids in every possible combination to yield peptides of a given length. Millions of chemical compounds theoretically can be synthesized through such combinatorial mixings of chemical building blocks. For example, one commentator observed that the systematic, combinatorial mixing of 100 interchangeable chemical building blocks results in the theoretical synthesis of 100 million tetrameric compounds or 10 billion pentameric compounds. (Gallop et al., "Applications of Combinatorial Technologies to Drug Discovery, Background and Peptide

Combinatorial Libraries," **Journal of Medicinal Chemistry**, Vol. 37, No. 9, 1233-1250 (1994). Other chemical libraries known to those in the art may also be used, including natural product libraries.

5 Once generated, combinatorial libraries can be screened for compounds that possess desirable biological properties. For example, compounds which may be useful as drugs or to develop drugs would likely have the ability to bind to the target protein identified, expressed and purified as discussed above. Further, if the identified target protein is an enzyme, candidate compounds would likely interfere with the enzymatic properties of the target protein. Any enzyme can be a target protein. For example, the  
10 enzymatic function of a target protein can be to serve as a protease, nuclease, phosphatase, dehydrogenase, transporter protein, transcriptional enzyme, and any other type of enzyme known or unknown. Thus, the present invention contemplates using the protein products described above to screen combinatorial chemical libraries.

Those in the art will appreciate that a number of techniques exist for characterizing  
15 target proteins in order to identify molecules useful for the discovery and development of therapeutics. For example, some techniques involve the generation and use of small peptides to probe and analyze target proteins both biochemically and genetically in order to identify and develop drug leads. Such techniques include the methods described in PCT publications No. WO9935494, WO9819162, WO9954728, the disclosures of which are  
20 incorporated herein by reference in their entireties.

In another example, the target protein is a serine protease and the substrate of the enzyme is known. The present example is directed towards the analysis of libraries of compounds to identify compounds that function as inhibitors of the target enzyme. First, a library of small molecules is generated using methods of combinatorial library formation  
25 well known in the art. U.S. Patent NOs. 5,463,564 and 5,574, 656, to Agrafiotis, et al., entitled "System and Method of Automatically Generating Chemical Compound with Desired Properties," are two such teachings. Then the library compounds are screened to identify library compounds that possess desired structural and functional properties. U.S. Patent No. 5,684,711 also discusses a method for screening libraries.

30 To illustrate the screening process, the combined target and chemical compounds of the library are exposed to and permitted to interact with the purified enzyme. A labeled substrate is added to the incubation. The label on the substrate is such that a detectable



signal is emitted from metabolized substrate molecules. The emission of this signal permits one to measure the effect of the combinatorial library compounds on the enzymatic activity of target enzymes. The characteristics of each library compound is encoded so that compounds demonstrating activity against the enzyme can be analyzed and features common to the various compounds identified can be isolated and combined into future iterations of libraries.

Once a library of compounds is screened, subsequent libraries are generated using those chemical building blocks that possess the features shown in the first round of screen to have activity against the target enzyme. Using this method, subsequent iterations of candidate compounds will possess more and more of those structural and functional features required to inhibit the function of the target enzyme, until a group of enzyme inhibitors with high specificity for the enzyme can be found. These compounds can then be further tested for their safety and efficacy as antibiotics for use in mammals.

It will be readily appreciated that this particular screening methodology is exemplary only. Other methods are well known to those skilled in the art. For example, a wide variety of screening techniques are known for a large number of naturally-occurring targets when the biochemical function of the target protein is known.

#### B. Cell Based Assays

Current cell-based assays used to identify or to characterize compounds for drug discovery and development frequently depend on detecting the ability of a test compound to inhibit the activity of a target molecule located within a cell or located on the surface of a cell. Most often such target molecules are proteins such as enzymes, receptors and the like. However, target molecules may also include other molecules such as DNAs, lipids, carbohydrates and RNAs including messenger RNAs, ribosomal RNAs, tRNAs and the like. A number of highly sensitive cell-based assay methods are available to those of skill in the art to detect binding and interaction of test compounds with specific target molecules. However, these methods are generally not highly effective when the test compound binds to or otherwise interacts with its target molecule with moderate or low affinity. In addition, the target molecule may not be readily accessible to a test compound in solution, such as when the target molecule is located inside the cell or within a cellular compartment such as the periplasm of a

bacterial cell. Thus, current cell-based assay methods are limited in that they are not effective in identifying or characterizing compounds that interact with their targets with moderate to low affinity or compounds that interact with targets that are not readily accessible.

5           Cell-based assay methods of the present invention have substantial advantages over current cell-based assays practiced in the art. These advantages derive from the use of sensitized cells in which the level or activity of a proliferation-required gene product (the target molecule) has been specifically reduced to the point where the presence or absence of its function becomes a rate-determining step for cellular proliferation. Bacterial, fungal, plant, or animal cells can all be used with the present method. Such sensitized cells become much more sensitive to compounds that are active against the affected target molecule. Thus, cell-based assays of the present invention are capable of detecting compounds exhibiting low or moderate potency against the target molecule of interest because such compounds are substantially more potent on sensitized cells than on non-sensitized cells. The affect may be such that a test compound may be two to several times more potent, at least 10 times more potent or even at least 100 times more potent when tested on the sensitized cells as compared to the non-sensitized cells.

20           Due in part to the increased appearance of antibiotic resistance in pathogenic microorganisms and to the significant side-effects associated with some currently used antibiotics, novel antibiotics acting at new targets are highly sought after in the art. Yet, another limitation in the current art related to cell-based assays is the problem of identifying hits against the same kinds of target molecules in the same limited set of biological pathways over and over again. This may occur when compounds acting at such new targets are discarded, ignored or fail to be detected because compounds acting at the "old" targets are encountered more frequently and are more potent than compounds acting at the new targets. As a result, the majority of antibiotics in use currently interact with a relatively small number of target molecules within an even more limited set of biological pathways.

30           The use of sensitized cells of the current invention provides a solution to the above problem in two ways. First, desired compounds acting at a target of interest, whether a new target or a previously known but poorly exploited target, can now be

detected above the "noise" of compounds acting at the "old" targets due to the specific and substantial increase in potency of such desired compounds when tested on the sensitized cells of the current invention. Second, the methods used to sensitize cells to compounds acting at a target of interest may also sensitize these cells to compounds acting at other target molecules within the same biological pathway. For example, expression of an antisense molecule to a gene encoding a ribosomal protein is expected to sensitize the cell to compounds acting at that ribosomal protein and may also sensitize the cells to compounds acting at any of the ribosomal components (proteins or rRNA) or even to compounds acting at any target which is part of the protein synthesis pathway. Thus an important advantage of the present invention is the ability to reveal new targets and pathways that were previously not readily accessible to drug discovery methods.

Sensitized cells of the present invention are prepared by reducing the activity or level of a target molecule. The target molecule may be a gene product, such as an RNA or polypeptide produced from the proliferation-required nucleic acids described herein. Alternatively, the target may be a gene product such as an RNA or polypeptide which is produced from a sequence within the same operon as the proliferation-required nucleic acids described herein. In addition, the target may be an RNA or polypeptide in the same biological pathway as the proliferation-required nucleic acids described herein. Such biological pathways include, but are not limited to, enzymatic, biochemical and metabolic pathways as well as pathways involved in the production of cellular structures such as the cell wall.

Current methods employed in the arts of medicinal and combinatorial chemistries are able to make use of structure-activity relationship information derived from testing compounds in various biological assays including direct binding assays and cell-based assays. Occasionally compounds are directly identified in such assays that are sufficiently potent to be developed as drugs. More often, initial hit compounds exhibit moderate or low potency. Once a hit compound is identified with low or moderate potency, directed libraries of compounds are synthesized and tested in order to identify more potent leads. Generally these directed libraries are combinatorial chemical libraries consisting of compounds with structures related to the hit compound but containing systematic variations including additions, subtractions and substitutions

of various structural features. When tested for activity against the target molecule, structural features are identified that either alone or in combination with other features enhance or reduce activity. This information is used to design subsequent directed libraries containing compounds with enhanced activity against the target molecule.

5 After one or several iterations of this process, compounds with substantially increased activity against the target molecule are identified and may be further developed as drugs. This process is facilitated by use of the sensitized cells of the present invention since compounds acting at the selected targets exhibit increased potency in such cell-based assays, thus; more compounds can now be characterized providing more useful  
10 information than would be obtained otherwise.

Thus, it is now possible using cell-based assays of the present invention to identify or characterize compounds that previously would not have been readily identified or characterized including compounds that act at targets that previously were not readily exploited using cell-based assays. The process of evolving potent drug leads  
15 from initial hit compounds is also substantially improved by the cell-based assays of the present invention because, for the same number of test compounds, more structure-function relationship information is likely to be revealed.

The method of sensitizing a cell entails selecting a suitable gene or operon. A suitable gene or operon is one whose expression is required for the proliferation of the  
20 cell to be sensitized. The next step is to introduce into the cells to be sensitized, an antisense RNA capable of hybridizing to the suitable gene or operon or to the RNA encoded by the suitable gene or operon. Introduction of the antisense RNA can be in the form of an expression vector in which antisense RNA is produced under the control of an inducible promoter. The amount of antisense RNA produced is limited by varying  
25 the inducer concentration to which the cell is exposed and thereby varying the activity of the promoter driving transcription of the antisense RNA. Thus, cells are sensitized by exposing them to an inducer concentration that results in a sub-lethal level of antisense RNA expression.

In one embodiment of the cell-based assays, the identified exogenous *E. coli*  
30 nucleotide sequences of the present invention are used to inhibit the production of a proliferation-required protein. Expression vectors producing antisense RNA against identified genes required for proliferation are used to limit the concentration of a

proliferation-required protein without severely inhibiting growth. To achieve that goal, a growth inhibition dose curve of inducer is calculated by plotting various doses of inducer against the corresponding growth inhibition caused by the antisense expression. From this curve, various percentages of antisense induced growth inhibition, from 1 to 100% can be determined. If the promoter contained in the expression vector contains a *lac* operator the transcription is regulated by *lac* repressor and expression from the promoter is inducible with IPTG. For example, the highest concentration of the inducer IPTG that does not reduce the growth rate (0% growth inhibition) can be predicted from the curve. Cellular proliferation can be monitored by growth medium turbidity via OD measurements. In another example, the concentration of inducer that reduces growth by 25% can be predicted from the curve. In still another example, a concentration of inducer that reduces growth by 50% can be calculated. Additional parameters such as colony forming units (cfu) can be used to measure cellular viability.

Cells to be assayed are exposed to the above-determined concentrations of inducer. The presence of the inducer at this sub-lethal concentration reduces the amount of the proliferation required gene product to the lowest amount in the cell that will support growth. Cells grown in the presence of this concentration of inducer are therefore specifically more sensitive to inhibitors of the proliferation-required protein or RNA of interest or to inhibitors of proteins or RNAs in the same biological pathway as the proliferation-required protein or RNA of interest but not to inhibitors of unrelated proteins or RNAs.

Cells pretreated with sub-inhibitory concentrations of inducer and thus containing a reduced amount of proliferation-required target gene product are then used to screen for compounds that reduce cell growth. The sub-lethal concentration of inducer may be any concentration consistent with the intended use of the assay to identify candidate compounds to which the cells are more sensitive. For example, the sub-lethal concentration of the inducer may be such that growth inhibition is at least about 5%, at least about 8%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60% at least about 75%, or more. Cells which are pre-sensitized using the preceding method are more sensitive to inhibitors of the target protein because these cells contain less target protein to inhibit than wild-type cells.

In another embodiment of the cell based assays of the present invention, the level or activity of a proliferation required gene product is reduced using a temperature sensitive ...mutation in the proliferation-required sequence and an antisense nucleic acid against the proliferation-required sequence. Growing the cells at an intermediate temperature  
5 between the permissive and restrictive temperatures of the temperature sensitive mutant where the mutation is in a proliferation-required gene produces cells with reduced activity of the proliferation-required gene product. The antisense RNA directed against the proliferation-required sequence further reduces the activity of the proliferation required gene product. Drugs that may not have been found using either the  
10 temperature sensitive mutation or the antisense nucleic acid alone may be identified by determining whether cells in which expression of the antisense nucleic acid has been induced and which are grown at a temperature between the permissive temperature and the restrictive temperature are substantially more sensitive to a test compound than cells in which expression of the antisense nucleic acid has not been induced and which are  
15 grown at a permissive temperature. Also drugs found previously from either the antisense nucleic acid alone or the temperature sensitive mutation alone may have a different sensitivity profile when used in cells combining the two approaches, and that sensitivity profile may indicate a more specific action of the drug in inhibiting one or more activities of the gene product.

20 Temperature sensitive mutations may be located at different sites within the gene and correspond to different domains of the protein. For example, the *dnaB* gene of *Escherichia coli* encodes the replication fork DNA helicase. DnaB has several domains, including domains for oligomerization, ATP hydrolysis, DNA binding, interaction with primase, interaction with DnaC, and interaction with DnaA [(Biswas,  
25 E.E. and Biswas, S.B. 1999. Mechanism and DnaB helicase of *Escherichia coli*: structural domains involved in ATP hydrolysis, DNA binding, and oligomerization. Biochem. **38**:10919-10928; Hiasa, H. and Marians, K.J. 1999. Initiation of bidirectional replication at the chromosomal origin is directed by the interaction between helicase and primase. J. Biol. Chem. **274**:27244-27248; San Martin, C., Rademacher, M.,  
30 Wolpensinger, B., Engel, A., Miles, C.S., Dixon, N.E., and Carazo, J.M. 1998. Three-dimensional reconstructions from cryoelectron microscopy images reveal an intimate complex between helicase DnaB and its loading partner DnaC. Structure **6**:501-9;

Sutton, M.D., Carr, K.M., Vicente, M., and Kaguni, J.M. 1998. *Escherichia coli* DnaA protein. The N-terminal domain and loading of DnaB helicase at the *E. coli* chromosomal. J. Biol. Chem. **273**:34255-62.), the disclosures of which are incorporated herein by reference in their entireties]. Temperature sensitive mutations in different domains of DnaB confer different phenotypes at the restrictive temperature, which include either an abrupt stop or slow stop in DNA replication with or without DNA breakdown (Wechsler, J.A. and Gross, J.D. 1971. *Escherichia coli* mutants temperature-sensitive for DNA synthesis. Mol. Gen. Genetics **113**:273-284, the disclosure of which is incorporated herein by reference in its entirety) and termination of growth or cell death. Combining the use of temperature sensitive mutations in the *dnaB* gene that cause cell death at the restrictive temperature with an antisense to the *dnaB* gene could lead to the discovery of very specific and effective inhibitors of one or a subset of activities exhibited by DnaB.

When screening for antimicrobial agents against a gene product required for proliferation, growth inhibition of cells containing a limiting amount of that proliferation-required gene product can be assayed. Growth inhibition can be measured by directly comparing the amount of growth, measured by the optical density of the growth medium, between an experimental sample and a control sample. Alternative methods for assaying cell proliferation include measuring green fluorescent protein (GFP) reporter construct emissions, various enzymatic activity assays, and other methods well known in the art.

It will be appreciated that the above method may be performed in solid phase, liquid phase or a combination of the two. For example, cells grown on nutrient agar containing the inducer of the antisense construct may be exposed to compounds spotted onto the agar surface. A compound's effect may be judged from the diameter of the resulting killing zone, the area around the compound application point in which cells do not grow. Multiple compounds may be transferred to agar plates and simultaneously tested using automated and semi-automated equipment including but not restricted to multi-channel pipettes (for example the Beckman Multimek) and multi-channel spotters (for example the Genomic Solutions Flexys). In this way multiple plates and thousands to millions of compounds may be tested per day.

5 The compounds may also be tested entirely in liquid phase using microtiter plates as described below. Liquid phase screening may be performed in microtiter plates containing 96, 384, 1536 or more wells per microtiter plate to screen multiple plates and thousands to millions of compounds per day. Automated and semi-automated equipment may be used for addition of reagents (for example cells and compounds) and determination of cell density.

#### EXAMPLE 9

10 The effectiveness of the above cell based assay was validated using constructs expressing antisense RNA to *E. coli* genes *rplL*, *rplJ*, and *rplW* encoding ribosomal proteins L7/L12, L10 and L23 respectively. These proteins are part of the protein synthesis apparatus of the cell and as such are required for proliferation. These constructs were used to test the effect of antisense expression on cell sensitivity to antibiotics known to bind to the ribosome and thereby inhibit protein synthesis. Constructs expressing antisense RNA to several other genes (*elaD*, *visC*, *yohH*, and  
15 *aptE/B*), the products of which are not involved in protein synthesis were used for comparison.

First expression vectors containing antisense constructs to either *rplW* or to *elaD* were introduced into separate *E. coli* cell populations. Vector introduction is a technique well known to those of ordinary skill in the art. The expression vectors of  
20 this example contain IPTG inducible promoters that drive the expression of the antisense RNA in the presence of the inducer. However, those skilled in the art will appreciate that other inducible promoters may also be used. Suitable expression vectors are also well known in the art. The *E. coli* antisense clones encoding ribosomal proteins L7/L12, L10 and L23 were used to test the effect of antisense expression on cell  
25 sensitivity to the antibiotics known to bind to these proteins. First, expression vectors containing antisense to either the genes encoding L7/L12 and L10 or L23 were introduced into separate *E. coli* cell populations.

The cell populations were exposed to a range of IPTG concentrations in liquid medium to obtain the growth inhibitory dose curve for each clone (Fig. 1). First, seed  
30 cultures were grown to a particular turbidity that is measured by the optical density (OD) of the growth solution. The OD of the solution is directly related to the number of bacterial cells contained therein. Subsequently, sixteen 200 ul liquid medium cultures



were grown in a 96 well microtiter plate at 37 C with a range of IPTG concentrations in duplicate two-fold serial dilutions from 1600  $\mu$ M to 12.5  $\mu$ M (final concentration). Additionally, control cells were grown in duplicate without IPTG. These cultures were started from equal amounts of cells derived from the same initial seed culture of a clone of interest. The cells were grown for up to 15 hours and the extent of growth was determined by measuring the optical density of the cultures at 600 nm. When the control culture reached mid-log phase the percent growth of the control for each of the IPTG containing cultures was plotted against the log concentrations of IPTG to produce a growth inhibitory dose response curve for the IPTG. The concentration of IPTG that inhibits cell growth to 50% ( $IC_{50}$ ) as compared to the 0 mM IPTG control (0% growth inhibition) was then calculated from the curve. Under these conditions, an amount of antisense RNA was produced that reduced the expression levels of *rplW* and *elaD* to a degree such that growth was inhibited by 50%.

Alternative methods of measuring growth are also contemplated. Examples of these methods include measurements of proteins, the expression of which is engineered into the cells being tested and can readily be measured. Examples of such proteins include green fluorescent protein (GFP) and various enzymes.

Cells were pretreated with the selected concentration of IPTG and then used to test the sensitivity of cell populations to tetracycline, erythromycin and other protein synthesis inhibitors. An example of a tetracycline dose response curve is shown in Figures 2A and 2B for the *rplW* and *elaD* genes, respectively. Cells were grown to log phase and then diluted into media alone or media containing IPTG at concentrations which give 20% and 50% growth inhibition as determined by IPTG dose response curves. After 2.5 hours, the cells were diluted to a final OD600 of 0.002 into 96 well plates containing (1) +/- IPTG at the same concentrations used for the 2.5 hour pre-incubation; and (2) serial two-fold dilutions of tetracycline such that the final concentrations of tetracycline range from 1  $\mu$ g/ml to 15.6 ng/ml and 0  $\mu$ g/ml. The 96 well plates were incubated at 37°C and the OD600 was read by a plate reader every 5 minutes for up to 15 hours. For each IPTG concentration and the no IPTG control, tetracycline dose response curves were determined when the control (absence of tetracycline) reached 0.1 OD600. To compare tetracycline sensitivity with and without IPTG, tetracycline  $IC_{50}$ s were determined from the dose response curves (Figs. 2A-B).

Cells with reduced levels of L23 (*rplW*) showed increased sensitivity to tetracycline (Fig. 2A) as compared to cells with reduced levels of *elaD* (Fig. 2B). Figure 3 shows a summary bar chart in which the ratios of tetracycline IC<sub>50</sub>s determined in the presence of IPTG which gives 50% growth inhibition versus tetracycline IC<sub>50</sub>s determined without IPTG (fold increase in tetracycline sensitivity) were plotted. Cells with reduced levels of either L7/L12 (genes *rplL*, *rplJ*) or L23 (*rplW*) showed increased sensitivity to tetracycline (Fig. 3). Cells expressing antisense to genes not known to be involved in protein synthesis (*atpB/E*, *visC*, *elaD*, *yohH*) did not show the same increased sensitivity to tetracycline, validating the specificity of this assay (Fig. 3).

In addition to the above, it has been observed in initial experiments that clones expressing antisense RNA to genes involved in protein synthesis (including genes encoding ribosomal proteins L7/L12 & L10, L7/L12 alone, L22, and L18, as well as genes encoding rRNA and Elongation Factor G) have increased sensitivity to the macrolide, erythromycin, whereas clones expressing antisense to the non-protein synthesis genes *elaD*, *atpB/E* and *visC* do not. Furthermore, the clone expressing antisense to *rplL* and *rplJ* does not show increased sensitivity to nalidixic acid and ofloxacin, antibiotics which do not inhibit protein synthesis.

The results with the ribosomal protein genes *rplL*, *rplJ*, and *rplW* as well as the initial results using various other antisense clones and antibiotics show that limiting the concentration of an antibiotic target makes cells more sensitive to the antimicrobial agents that specifically interact with that protein. The results also show that these cells are sensitized to antimicrobial agents that inhibit the overall function in which the protein target is involved but are not sensitized to antimicrobial agents that inhibit other functions.

The cell based assay described above may also be used to identify the biological pathway in which a proliferation-required nucleic acid or its gene product lies. In such methods, cells expressing a sub-lethal level of antisense to a target proliferation-required nucleic acid and control cells in which expression of the antisense has not been induced are contacted with a panel of antibiotics known to act in various pathways. If the antibiotic acts in the pathway in which the target proliferation-required nucleic acid or its gene product lies, cells in which expression of the antisense has been induced will

be more sensitive to the antibiotic than cells in which expression of the antisense has not been induced.

As a control, the results of the assay may be confirmed by contacting a panel of cells expressing antisense nucleic acids to many different proliferation-required genes including the target proliferation-required gene. If the antibiotic is acting specifically, heightened sensitivity to the antibiotic will be observed only in the cells expressing antisense to a target proliferation-required gene (or cells expressing antisense to other proliferation-required genes in the same pathway as the target proliferation-required gene) but will not be observed generally in all cells expressing antisense to proliferation-required genes.

Similarly, the above method may be used to determine the pathway on which a test antibiotic acts. A panel of cells, each of which expresses antisense to a proliferation-required nucleic acid in a known pathway, is contacted with a compound for which it is desired to determine the pathway on which it acts. The sensitivity of the panel of cells to the test compound is determined in cells in which expression of the antisense has been induced and in control cells in which expression of the antisense has not been induced. If the test antibiotic acts on the pathway on which an antisense nucleic acid acts, cells in which expression of the antisense has been induced will be more sensitive to the antibiotic than cells in which expression of the antisense has not been induced. In addition, control cells in which expression of antisense to proliferation-required genes in other pathways has been induced will not exhibit heightened sensitivity to the antibiotic. In this way, the pathway on which the test antibiotic acts may be determined.

The Example below provides one method for performing such assays.

#### **EXAMPLE 10**

##### Identification of the Pathway in which a Proliferation-Required Gene Lies or the Pathway on which an Antibiotic Acts

###### A. Preparation of Bacterial Stocks for Assay

To provide a consistent source of cells to screen, frozen stocks of host bacteria containing the desired antisense construct are prepared using standard microbiological techniques. For example, a single clone of the organism can be isolated by streaking out a sample of the original stock onto an agar plate containing nutrients for cell growth and

an antibiotic for which the antisense construct contains a gene which confers resistance. After overnight growth an isolated colony is picked from the plate with a sterile needle and transferred to an appropriate liquid growth media containing the antibiotic required for maintenance of the plasmid. The cells are incubated at 30°C to 37°C with vigorous shaking for 4 to 6 hours to yield a culture in exponential growth. Sterile glycerol is added to 15% (volume to volume) and 100 µL to 500 µL aliquots are distributed into sterile cryotubes, snap frozen in liquid nitrogen, and stored at -80°C for future assays.

#### B. Growth of Bacteria for Use in the Assay

A day prior to an assay, a stock vial is removed from the freezer, rapidly thawed (37°C water bath) and a loop of culture is streaked out on an agar plate containing nutrients for cell growth and an antibiotic to which the antisense construct confers resistance. After overnight growth at 37°C, ten randomly chosen, isolated colonies are transferred from the plate (sterile inoculum loop) to a sterile tube containing 5 mL of LB medium containing the antibiotic to which the antisense vector confers resistance. After vigorous mixing to form a homogeneous cell suspension, the optical density of the suspension is measured at 600 nm (OD600) and if necessary an aliquot of the suspension is diluted into a second tube of 5 mL. sterile, LB medium plus antibiotic to achieve an  $OD_{600} \leq 0.02$  absorbance units. The culture is then incubated at 37° C for 1-2 hrs with shaking until the OD600 reaches OD 0.2 – 0.3. At this point the cells are ready to be used in the assay.

#### C. Selection of Media to be Used in Assay

Two fold dilution series of the inducer are generated in culture media containing the appropriate antibiotic for maintenance of the antisense construct. Several media are tested side by side and three to four wells are used to evaluate the effects of the inducer at each concentration in each media. For example, M9 minimal media, LB broth, TBD broth and Muller-Hinton media may be tested with the inducer IPTG at the following concentrations, 50 µM, 100 µM, 200 µM, 400 µM, 600 µM, 800 µM and 1000 µM. Equal volumes of test media-inducer and cells are added to the wells of a 384 well microtiter plate and mixed. The cells are prepared as described above and diluted 1:100 in the appropriate media containing the test antibiotic immediately prior to addition to the microtiter plate wells. For a control, cells are also added to several wells of each media that do not contain inducer, for example 0 µM IPTG. Cell growth is monitored

continuously by incubation at 37°C in a microtiter plate reader monitoring the OD600 of the wells over an 18-hour period. The percent inhibition of growth produced by each concentration of inducer is calculated by comparing the rates of logarithmic growth against that exhibited by cells growing in media without inducer. The medium yielding  
5 greatest sensitivity to inducer is selected for use in the assays described below.

D. Measurement of Test Antibiotic Sensitivity in the Absence of Antisense Construct Induction

Two-fold dilution series of antibiotics of known mechanism of action are generated in the culture media selected for further assay development that has been  
10 supplemented with the antibiotic used to maintain the construct. A panel of test antibiotics known to act on different pathways is tested side by side with three to four wells being used to evaluate the effect of a test antibiotic on cell growth at each concentration. Equal volumes of test antibiotic and cells are added to the wells of a 384 well microtiter plate and mixed. Cells are prepared as described above using the media  
15 selected for assay development supplemented with the antibiotic required to maintain the antisense construct and are diluted 1:100 in identical media immediately prior to addition to the microtiter plate wells. For a control, cells are also added to several wells that contain the solvent used to dissolve the antibiotics but no antibiotic. Cell growth is monitored continuously by incubation at 37°C in a microtiter plate reader monitoring  
20 the OD600 of the wells over an 18-hour period. The percent inhibition of growth produced by each concentration of antibiotic is calculated by comparing the rates of logarithmic growth against that exhibited by cells growing in media without antibiotic. A plot of percent inhibition against log[antibiotic concentration] allows extrapolation of an IC<sub>50</sub> value for each antibiotic.

25 E. Measurement of Test Antibiotic Sensitivity in the Presence of Antisense Construct Inducer

The culture media selected for use in the assay is supplemented with inducer at concentrations shown to inhibit cell growth by 50 and 80% as described above and the antibiotic used to maintain the construct. Two fold dilution series of the panel of test  
30 antibiotics used above are generated in each of these media. Several antibiotics are tested side by side with three to four wells being used to evaluate the effects of an antibiotic on cell growth at each concentration, in each media. Equal volumes of test

antibiotic and cells are added to the wells of a 384 well microtiter plate and mixed. Cells are prepared as described above using the media selected for use in the assay supplemented with the antibiotic required to maintain the antisense construct. The cells are diluted 1:100 into two 50 mL aliquots of identical media containing concentrations of inducer that have been shown to inhibit cell growth by 50% and 80 % respectively and incubated at 37°C with shaking for 2.5 hours. Immediately prior to addition to the microtiter plate wells, the cultures are adjusted to an appropriate OD<sub>600</sub> (typically 0.002) by dilution into warm (37°C) sterile media supplemented with identical concentrations of the inducer and antibiotic used to maintain the antisense construct. For a control, cells are also added to several wells that contain solvent used to dissolve test antibiotics but which contain no antibiotic. Cell growth is monitored continuously by incubation at 37°C in a microtiter plate reader monitoring the OD<sub>600</sub> of the wells over an 18-hour period. The percent inhibition of growth produced by each concentration of antibiotic is calculated by comparing the rates of logarithmic growth against that exhibited by cells growing in media without antibiotic. A plot of percent inhibition against log[antibiotic concentration] allows extrapolation of an IC<sub>50</sub> value for each antibiotic.

#### F. Determining the Specificity of the Test Antibiotics

A comparison of the IC<sub>50</sub>s generated by antibiotics of known mechanism of action under antisense induced and non-induced conditions allows the pathway in which a proliferation-required nucleic acid lies to be identified. If cells expressing an antisense nucleic acid against a proliferation-required gene are selectively sensitive to an antibiotic acting via a particular pathway, then the gene against which the antisense acts is involved in the pathway in which the antibiotic acts.

#### G. Identification of Pathway in which a Test Antibiotic Acts

As discussed above, the cell based assay may also be used to determine the pathway against which a test antibiotic acts. In such an analysis, the pathways against which each member of a panel of antisense nucleic acids acts are identified as described above. A panel of cells, each containing an inducible antisense vector against a gene in a known proliferation-required pathway, is contacted with a test antibiotic for which it is desired to determine the pathway on which it acts under inducing an non-inducing conditions. If heightened sensitivity is observed in induced cells expressing antisense against a gene in a particular pathway but not in induced cells expressing antisense

against genes in other pathways, then the test antibiotic acts against the pathway for which heightened sensitivity was observed.

One skilled in the art will appreciate that further optimization of the assay conditions, such as the concentration of inducer used to induce antisense expression and/or the growth conditions used for the assay (for example incubation temperature and media components) may further increase the selectivity and/or magnitude of the antibiotic sensitization exhibited.

The following example confirms the effectiveness of the methods described above.

### EXAMPLE 11

#### Identification of the Pathway in which a Proliferation-Required Gene Lies

Antibiotics of various chemical classes and modes of action were purchased from Sigma Chemicals (St. Louis, MO). Stock solutions were prepared by dissolving each antibiotic in an appropriate aqueous solution based on information provided by the manufacturer. The final working solution of each antibiotic contained no more than 0.2% (w/v) of any organic solvent. To determine their potency against a bacterial strain engineered for expression of an antisense against a proliferation-required 50S ribosomal protein, each antibiotic was serially diluted two or three fold in growth medium supplemented with the appropriate antibiotic for maintenance of the anti-sense construct. At least ten dilutions were prepared for each antibiotic. 25  $\mu$ L aliquots of each dilution were transferred to discrete wells of a 384-well microplate (the assay plate) using a multi-channel pipette. Quadruplicate wells were used for each dilution of an antibiotic under each treatment condition (plus and minus inducer). Each assay plate contained twenty wells for cell growth controls (growth media replacing antibiotic), ten wells for each treatment (plus and minus inducer, in this example IPTG). Assay plates were usually divided into the two treatments: half the plate containing induced cells and an appropriate concentrations of inducer (in this example IPTG) to maintain the state of induction, the other half containing non-induced cells in the absence of IPTG.

Cells for the assay were prepared as follows. Bacterial cells containing a construct, from which expression of antisense nucleic acid against rplL and rplJ, which encode proliferation-required 50S ribosomal subunit proteins, is inducible in the presence of IPTG, were grown into exponential growth ( $OD_{600}$  0.2 to 0.3) and then

diluted 1:100 into fresh media containing either 400  $\mu$ M or 0  $\mu$ M inducer (IPTG). These cultures were incubated at 37° C for 2.5 hr. After a 2.5 hr incubation, induced and non-induced cells were respectively diluted into an assay medium at a final OD<sub>600</sub> value of 0.0004. The medium contained an appropriate concentration of the antibiotic for the maintenance of the anti-sense construct. In addition, the medium used to dilute induced cells was supplemented with 800  $\mu$ M IPTG so that addition to the assay plate would result in a final IPTG concentration of 400  $\mu$ M. Induced and non-induced cell suspensions were dispensed (25  $\mu$ l/well) into the appropriate wells of the assay plate as discussed previously. The plate was then loaded into a plate reader, incubated at constant temperature, and cell growth was monitored in each well by the measurement of light scattering at 595 nm. Growth was monitored every 5 minutes until the cell culture attained a stationary growth phase. For each concentration of antibiotic, a percentage inhibition of growth was calculated at the time point corresponding to mid-exponential growth for the associated control wells (no antibiotic, plus or minus IPTG). For each antibiotic and condition (plus or minus IPTG), a plot of percent inhibition versus log of antibiotic concentration was generated and the IC<sub>50</sub> determined. A comparison of the IC<sub>50</sub> for each antibiotic in the presence and absence of IPTG revealed whether induction of the antisense construct sensitized the cell to the mechanism of action exhibited by the antibiotic. Cells which exhibited a significant (standard statistical analysis) numerical decrease in the IC<sub>50</sub> value in the presence of inducer were considered to have an increased sensitivity to the test antibiotic.

The results are provided in the table below, which lists the classes and names of the antibiotics used in the analysis, the targets of the antibiotics, the IC<sub>50</sub> in the absence of IPTG, the IC<sub>50</sub> in the presence of IPTG, the concentration units for the IC<sub>50</sub>s, the fold increase in IC<sub>50</sub> in the presence of IPTG, and whether increased sensitivity was observed in the presence of IPTG.



**TABLE IV**  
**Effect of Expression of Antisense RNA to rplL and rplJ on Antibiotic Sensitivity**

ANTIBIOTIC CLASS /Names	TARGET	IC50 (-IPTG)	IC50 (+IPTG)	Conc. Unit	Fold Increase in Sensitivity	Sensitivity Increased?
<b>PROTEIN SYNTHESIS INHIBITOR</b>						
<b>ANTIBIOTICS</b>						
<b>AMINOGLYCOSIDES</b>						
Gentamicin	30S ribosome function	2715	19.19	ng/ml	141	Yes
Streptomycin	30S ribosome function	11280	161	ng/ml	70	Yes
Spectinomycin	30S ribosome function	18050	<156	ng/ml		Yes
Tobramycin	30S ribosome function	3594	70.58	ng/ml	51	Yes
<b>MACROLIDES</b>						
Erythromycin	50S ribosome function	7467	187	ng/ml	40	Yes
<b>AROMATIC POYKETIDES</b>						
Tetracycline	30S ribosome function	199.7	1.83	ng/ml	109	Yes
Minocycline	30S ribosome function	668.4	3.897	ng/ml	172	Yes
Doxycycline	30S ribosome function	413.1	27.81	ng/ml	15	Yes
<b>OTHER PROTEIN SYNTHESIS INHIBITORS</b>						
Fusidic acid	Elongation Factor G function	59990	641	ng/ml	94	Yes
Chloramphenicol	30S ribosome function	465.4	1.516	ng/ml	307	Yes
Lincomycin	50S ribosome function	47150	324.2	ng/ml	145	Yes
<b>OTHER ANTIBIOTIC MECHANISMS</b>						
<b>B-LACTAMS</b>						
Cefoxitin	Cell wall biosynthesis	2782	2484	ng/ml	1	No
Cefotaxime	Cell wall biosynthesis	24.3	24.16	ng/ml	1	No
<b>DNA SYNTHESIS INHIBITORS</b>						
Nalidixic acid	DNA Gyrase activity	6973	6025	ng/ml	1	No
Ofloxacin	DNA Gyrase activity	49.61	45.89	ng/ml	1	No

OTHER	Cell membrane function	4077	4677	mg/ml	1	No
Bacitracin	Dihydrofolate Reductase activity	128.9	181.97	ng/ml	1	No
Trimethoprim	Cell wall biosynthesis	145400	72550	ng/ml	2	No
Vancomycin						

The above results demonstrate that induction of an antisense RNA to genes encoding 50S ribosomal subunit proteins results in a selective and highly significant sensitization of cells to antibiotics that inhibit ribosomal function and protein synthesis. The above results further demonstrate that induction of an antisense construct to an essential gene sensitizes an organism to compounds that interfere with that gene products' biological role. This sensitization is restricted to compounds that interfere with pathways associated with the targeted gene and it's product.

Assays utilizing antisense constructs to essential genes can be used to identify compounds that specifically interfere with the activity of multiple targets in a pathway. Such constructs can be used to simultaneously screen a sample against multiple targets in one pathway in one reaction (Combinatorial HTS).

Furthermore, as discussed above, panels of antisense construct containing cells may be used to characterize the point of intervention of any compound affecting an essential biological pathway including antibiotics with no known mechanism of action.

Another embodiment of the present invention is a method for determining the pathway against which a test antibiotic compound is active in which the activity of target proteins or nucleic acids involved in proliferation-required pathways is reduced by contacting cells with a sublethal concentration of a known antibiotic which acts against the target protein or nucleic acid. In one embodiment, the target protein or nucleic acid is a target protein or nucleic acid corresponding to a proliferation-required nucleic acid identified using the methods described above. The method is similar to those described above for determining which pathway a test antibiotic acts against except that rather than reducing the activity or level of a proliferation-required gene product using a sublethal level of antisense to a proliferation-required nucleic acid, the activity or level of the proliferation-required gene product is reduced using sublethal level of a known antibiotic which acts against the proliferation required gene product.

Interactions between drugs which affect the same biological pathway has been described in the literature. For example, Mecillinam (Amdinocillin) binds to and inactivates the penicillin binding protein 2 (PBP2, product of the *mrdA* in *E. coli*). This antibiotic inteacts with other antibiotics that inhibit PBP2 as well as antibiotics that inhibit other penicillin binding proteins such as PBP3 [(Gutmann, L., Vincent, S., Billot-Klein, D., Acar, J.F., Mrena, E., and Williamson, R. (1986) Involvement of

penicillin-binding protein 2 with other penicillin-binding proteins in lysis of *Escherichia coli* by some beta-lactam antibiotics alone and in synergistic lytic effect of amdinocillin (mecillinam). Antimicrobial Agents & Chemotherapy, 30:906-912), the disclosure of which is incorporated herein by reference in its entirety]. Interactions  
5 between drugs could, therefore, involve two drugs that inhibit the same target protein or nucleic acid or inhibit different proteins or nucleic acids in the same pathway [(Fukuoka, T., Domon, H., Kakuta, M., Ishii, C., Hirasawa, A., Utsui, Y., Ohya, S., and Yasuda, H. (1997) Combination effect between panipenem and vancomycin on highly methicillin-resistant *Staphylococcus aureus*. Japan. J. Antibio. 50:411-419; Smith, C.E.,  
10 Foleno, B.E., Barrett, J.F., and Frosc, M.B. (1997) Assessment of the synergistic interactions of levofloxacin and ampicillin against *Enterococcus faecium* by the checkerboard agar dilution and time-kill methods. Diagnos. Microbiol. Infect. Disease 27:85-92; den Hollander, J.G., Horrevorts, A.M., van Goor, M.L., Verbrugh, H.A., and Mouton, J.W. (1997) Synergism between tobramycin and ceftazidime against a resistant  
15 *Pseudomonas aeruginosa* strain, tested in an in vitro pharmacokinetic model. Antimicrobial Agents & Chemotherapy. 41:95-110), the disclosure of all of which are incorporated herein by reference in their entirety].

Two drugs may interact even though they inhibit different targets. For example, the proton pump inhibitor, Omeprazole, and the antibiotic, Amoxycillin, two synergistic  
20 compounds acting together, can cure *Helicobacter pylori* infection [( Gabryelewicz, A., Laszewicz, W., Dzieniszewski, J., Ciok, J., Marlicz, K., Bielecki, D., Popiela, T., Legutko, J., Knapik, Z., Poniewierka, E. (1997) Multicenter evaluation of dual-therapy (omeprazol and amoxycillin) for *Helicobacter pylori*-associated duodenal and gastric ulcer (two years of the observation). J. Physiol. Pharmacol. 48 Suppl 4:93-105), the  
25 disclosure of which is incorporated herein by reference in its entirety].

The growth inhibition from the sublethal concentration of the known antibiotic may be at least about 5%, at least about 8%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, or at least about 75%, or more.

30 Alternatively, the sublethal concentration of the known antibiotic may be determined by measuring the activity of the target proliferation-required gene product rather than by measuring growth inhibition.

Cells are contacted with a combination of each member of a panel of known antibiotics at a sublethal level and varying concentrations of the test antibiotic. As a control, the cells are contacted with varying concentrations of the test antibiotic alone. The  $IC_{50}$  of the test antibiotic in the presence and absence of the known antibiotic is determined. If the  $IC_{50}$ s in the presence and absence of the known drug are substantially similar, then the test drug and the known drug act on different pathways. If the  $IC_{50}$ s are substantially different, then the test drug and the known drug act on the same pathway.

Another embodiment of the present invention is a method for identifying a candidate compound for use as an antibiotic in which the activity of target proteins or nucleic acids involved in proliferation-required pathways is reduced by contacting cells with a sublethal concentration of a known antibiotic which acts against the target protein or nucleic acid. In one embodiment, the target protein or nucleic acid is a target protein or nucleic acid corresponding to a proliferation-required nucleic acid identified using the methods described above. The method is similar to those described above for identifying candidate compounds for use as antibiotics except that rather than reducing the activity or level of a proliferation-required gene product using a sublethal level of antisense to a proliferation-required nucleic acid, the activity or level of the proliferation-required gene product is reduced using a sublethal level of a known antibiotic which acts against the proliferation required gene product.

The growth inhibition from the sublethal concentration of the known antibiotic may be at least about 5%, at least about 8%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, or at least about 75%, or more.

Alternatively, the sublethal concentration of the known antibiotic may be determined by measuring the activity of the target proliferation-required gene product rather than by measuring growth inhibition.

In order to characterize test compounds of interest, cells are contacted with a panel of known antibiotics at a sublethal level and one or more concentrations of the test compound. As a control, the cells are contacted with the same concentrations of the test compound alone. The  $IC_{50}$  of the test compound in the presence and absence of the known antibiotic is determined. If the  $IC_{50}$  of the test compound is substantially

different in the presence and absence of the known drug then the test compound is a good candidate for use as an antibiotic. As discussed above, once a candidate compound is identified using the above methods its structure may be optimized using standard techniques such as combinatorial chemistry.

- 5 Representative known antibiotics which may be used in each of the above methods are provided in the table below. However, it will be appreciated that other antibiotics may also be used.

ANTIBIOTIC	INHIBITS/TARGET	RESISTANT MUTANTS
<b>Inhibitors of Transcription</b>		
Rifamycin, 1959 Rifampicin	Inhibits initiation of transcription/ $\beta$ -subunit RNA polymerase, <i>rpoB</i>	<i>rpoB</i> , <i>crp</i> , <i>cyaA</i>
Rifabutin Rifaximin	Accelerates transcription chain termination/ $\beta$ -subunit RNA polymerase	<i>rpoB</i>
Streptolydigin	an acyclic ansamycin, inhibits RNA polymerase	<i>rpoB</i>
Streptovaricin	Intercalates between 2 successive G-C pairs, <i>rpoB</i> , inhibits RNA synthesis	<i>pIdA</i>
Actinomycin D+EDTA		
<b>Inhibitors of Nucleic Acid Metabolism</b>		
Quinolones, 1962 Nalidixic acid Oxolinic acid	$\alpha$ subunit gyrase and/or topoisomerase IV, <i>gyrA</i>	<i>gyrA</i> , <i>AorB</i> , <i>icd</i> , <i>sloB</i>
Fluoroquinolones	$\alpha$ subunit gyrase, <i>gyrA</i> and/or topoisomerase IV (probable target in Staph)	<i>gyrA</i>
Ciprofloxacin, 1983		<i>norA</i> (efflux in Staph)
Norfloxacin		<i>hlpQ</i>
Coumerins Novobiocin	Inhibits ATPase activity of $\beta$ -subunit gyrase, <i>gyrB</i>	<i>gyrB</i> , <i>cysB</i> , <i>cysE</i> , <i>nov</i> , <i>ompA</i>
Coumermycin	Inhibits ATPase activity of $\beta$ -subunit gyrase, <i>gyrB</i>	<i>gyrB</i> , <i>hisW</i>
Albicidin	DNA synthesis	<i>tsx</i> (nucleoside channel)
Metronidazole	Causes single-strand breaks in DNA	<i>nar</i>
<b>Inhibitors of Metabolic Pathways</b>		
Sulfonamides, 1932	blocks synthesis of dihydrofolate, dihydro-pterolate synthesis, <i>folP</i>	<i>folP</i> , <i>gpt</i> , <i>pabA</i> , <i>pabB</i> , <i>pabC</i>
Sulfanilamide		
Trimethoprim, 1962	Inhibits dihydrofolate reductase, <i>folA</i>	<i>folA</i> , <i>thyA</i>
Showdomycin	Nucleoside analogue capable of alkylating sulfhydryl groups, inhibitor of thymidylate synthetase	<i>nupC</i> , <i>pnp</i>
Thiolactomycin	type II fatty acid synthase inhibitor	<i>emrB</i>
Psicofuranine	Adenosine glycoside antibiotic, target is GMP synthetase	<i>fadB</i> , <i>emrB</i> due to gene dosage
Triclosan	Inhibits fatty acid synthesis	<i>guaA</i> , <i>B</i>
Diazoborines Isoniazid,	heterocyclic, contains boron, inhibit fatty acid synthesis, enoyl-ACP reductase, <i>fabI</i>	<i>fabI</i> ( <i>envM</i> )
Ethionamide		<i>fabI</i> ( <i>envM</i> )

# **Inhibitors of Translation**

Phenylpropanoids Chloramphenicol, 1947	Binds to ribosomal peptidyl transfer center preventing peptide translocation/ binds to S6, L3, L6, L14, L16, L25, L26, L27, but preferentially to L16	<i>rrn, cmlA, marA, ompF, ompR</i>
Tetracyclines, 1948, type II polyketides Minocycline Doxycycline	Binding to 30S ribosomal subunit, "A" site on 30S subunit, blocks peptide elongation, strongest binding to S7	<i>clmA (cmr), mar, ompF</i>
Macrolides (type I polyketides) Erythromycin, 1950 Carbomycin, Spiramycin etc	Binding to 50 S ribosomal subunit, 23S rRNA, blocks peptide translocation, L15, L4, L12	<i>rrn, rplC, rplD, rplV, mac</i>
Aminoglycosides Streptomycin, 1944 Neomycin  Spectinomycin Kanamycin  Kasugamycin  Gentamicin, 1963 Amikacin Paromycin	Irreversible binding to 30S ribosomal subunit, prevents translation or causes mistranslation of mRNA/16S rRNA	<i>rpsL, strC, M, ubiF atpA-E, ecfB. hemAC, D, E, G, topA, rpsC, D, E, rrn, spcB atpA-atpE, cpxA, ecfB, hemA, B, L, topA ksgA, B, C, D, rplB, K, rpsI, N, M, R rplF, ubiF cpxA rpsL</i>
Lincosamides Lincomycin, 1955 Clindamycin	Binding to 50 S ribosomal subunit, blocks peptide translocation	<i>linB, rplN, O, rpsG</i>
Streptogramins Virginiamycin, 1955 Pristinamycin Synercid: quinupristin /dalfopristin	2 components, Streptogramins A&B, bind to the 50S ribosomal subunit blocking peptide translocation and peptide bond formation	
Fusidanes Fusidic Acid Kirromycin (Mocimycin)	Inhibition of elongation factor G (EF-G) prevents peptide translocation Inhibition of elongation factor TU (EF-Tu), prevents peptide bond formation	<i>fusA ufuA, B</i>
Pulvomycin Thiopeptin	Binds to and inhibits EF-TU Sulfur-containing antibiotic, inhibits protein synthesis, EF-G	<i>rplE</i>
Tiamulin Negamycin	Inhibits protein synthesis Inhibits termination process of protein synthesis	<i>rplC, rplD prfB</i>
Oxazolidinones Linezolid Isoniazid	23S rRNA	
Nitrofurantoin	Inhibits protein synthesis. nitroreductases convert nitrofurantoin to highly reactive electrophilic intermediates which attack bacterial ribosomal proteins non-specifically	<i>pdx nfnA, B</i>
Pseudomonic Acids Mupirocin (Bactroban)	Inhibition of isoleucyl tRNA synthetase-used for Staph, topical cream, nasal spray	<i>ileS</i>

Indolmycin  
Viomycin

Inhibits tryptophanyl-tRNA synthetase

*trpS*  
*rrmA* (23S rRNA  
methyltransferase;  
mutant has slow  
growth rate, slow  
chain elongation rate,  
and viomycin  
resistance)

Thiopeptides  
Thiostrepton  
Micrococcin

Binds to L11-23S RNA complex  
Inhibits GTP hydrolysis by EF-G  
Stimulates GTP hydrolysis by EF-G

#### Inhibitors of Cell Walls/Membranes

$\beta$ -lactams

Penicillin, 1929 Ampicillin

Methicillin, 1960

Inhibition of one or more cell wall  
transpeptidases, endopeptidases, and  
glycosidases (PBPs), of the 12 PBPs only 2  
are essential: *mrda* (PBP2) and *fisI* (*pbpB*,  
PBP3)

*ampC*, *ampD*, *ampE*,  
*envZ*, *galU*, *hipA*,  
*hipQ*, *ompC*, *ompF*,  
*ompR*, *ptsI*, *rfa*,  
*tolD*, *tolE*

Cephalosporins, 1962

Mecillinam (amdinocillin)

Binds to and inactivates PBP2 (*mrda*)  
Inactivates PBP3 (*fisI*)

*tonB*  
*alaS*, *argS*, *crp*, *cyaA*,  
*envB*, *mrda*, *B*,  
*mreB*, *C*, *D*

Aztreonam (Furazlocillin)

Bacilysin, Tetaine

Glycopeptides Vancomycin, 1955

Dipeptide, inhib glucosamine synthase  
Inhib G+ cell wall syn, binds to terminal  
D-ala-D-ala of pentapeptide,  
Prevents dephosphorylation and  
regeneration of lipid carrier  
Disrupts multiple aspects of membrane  
function, including peptidoglycan  
synthesis, lipoteichoic acid synthesis, and  
the bacterial membrane potential

*dppA*

Polypeptides Bacitracin

Cyclic lipopeptide Daptomycin,  
1980

Surfactant action disrupts cell membrane  
lipids, binds lipid A moiety of LPS  
Analogue of P-enolpyruvate, inhibits 1<sup>st</sup>  
step in peptidoglycan synthesis - UDP-N-  
acetylglucosamine enolpyruvyl  
transferase, *murA*. Also acts as  
Immunosuppressant

*rfa*

Cycloserine

Alafosfalin

Prevents formation of D-ala dimer,  
inhibits D-ala ligase, *ddlA*, *B*  
phosphonodipeptide, cell wall synthesis  
inhibitor, potentiator of  $\beta$ -lactams

*pmrA*

*murA*, *crp*, *cyaA*, *glpT*,  
*hipA*, *ptsI*, *uhpT*

*hipA*, *cycA*

*pepA*, *tpp*

#### Inhibitors of Protein Processing/Transport

Globomycin

Inhibits signal peptidase II (cleaves  
prolipoproteins subsequent to lipid  
modification, *lspA*

*lpp*, *dnaE*



## EXAMPLE 12

### Transfer of Exogenous Nucleic Acid Sequences to other Bacterial Species Using the *E. coli* Expression Vectors or Expression Vectors Functional in Bacterial Species other than *E. coli*.

5           The above methods were validated using antisense nucleic acids which inhibit the growth of *E. coli* which were identified using methods similar to those described above. Expression vectors which inhibited growth of *E. coli* upon induction of antisense RNA expression with IPTG were transformed directly into *Enterobacter cloacae*, *Klebsiella pneumonia* or *Salmonella typhimurium*. The transformed cells were  
10 then assayed for growth inhibition according to the method of Example 1. After growth in liquid culture, cells were plated at various serial dilutions and a score determined by calculating the log difference in growth for INDUCED vs. UNINDUCED antisense RNA expression as determined by the maximum 10 fold dilution at which a colony was observed. The results of these experiments are listed below in Table VI. If there was  
15 no effect of antisense RNA expression in an organism, the clone is minus in Table VI. In contrast, a positive in Table VI means that at least 10 fold more cells were required to observe a colony on the induced plate than on the non-induced plate under the conditions used and in that organism.

20           Sixteen of the constructs were found to inhibit growth in all the organisms tested upon induction of antisense RNA expression with IPTG. Those skilled in the art will appreciate that a negative result in a heterologous organism does not mean that that organism is missing that gene nor does it mean that the gene is unessential. However, a positive result means that the heterologous organism contains a homologous gene which is required for proliferation of that organism. The homologous gene may be obtained  
25 using the methods described herein. Those cells that are inhibited by antisense may be used in cell based assays as described herein for the identification and characterization of compounds in order to develop antibiotics effective in these organisms. Those skilled in the art will appreciate that an antisense molecule which works in the organism from which it was obtained will not always work in a heterologous organism.

30

TABLE VI  
Sensitivity of Other Microorganisms to Antisense Nucleic Acids That Inhibit  
Proliferation in *E. coli*

5

Mol. No.	<i>S. typhimurium</i>	<i>E. cloacae</i>	<i>K. pneumoniae</i>
EcXA001	+	+	-
EcXA004	-	-	-
EcXA005	+	+	+
EcXA006	-	-	-
EcXA007	-	+	-
EcXA008	+	-	+
EcXA010	+	+	+
EcXA011	-	+	-
EcXA012	-	+	-
EcXA013	+	+	+
EcXA014	+	+	-
EcXA015	-	+	+
EcXA016	+	+	+
EcXA017	+	+	+
EcXA018	+	+	+
EcXA019	+	+	+
EcXA020	+	+	+
EcXA021	+	+	+
EcXA023	+	+	+
EcXA024	+	-	+
EcXA025	-	-	-

Mol. No.	<i>S. typhimurium</i>	<i>E. cloacae</i>	<i>K. pneumoniae</i>
EcXA026	+	+	-
EcXA027	+	+	+
EcXA028	+	-	-
EcXA029	-	-	-
EcXA030	+	+	+
EcXA031	+	-	-
EcXA032	+	-	-
EcXA033	+	+	+
EcXA034	+	+	+
EcXA035	-	-	-
EcXA036	+	-	+
EcXA037	-	+	-
EcXA038	+	+	-
EcXA039	+	-	-
EcXA041	+	+	+
EcXA042	-	+	+
EcXA044	-	-	-
EcXA045	-	+	-
EcXA046	-	-	-
EcXA047	+	+	-
EcXA048	-	-	-
EcXA049	+	-	-
EcXA050	-	-	-

Mol. No.	<i>S. typhimurium</i>	<i>E. cloacae</i>	<i>K. pneumoniae</i>
EcXA051	+	-	-
EcXA052	+	-	-
EcXA053	+	+	+
EcXA054	-	-	+
EcXA055	+	-	-

### EXAMPLE 13

#### Use of Identified Exogenous Nucleic Acid Sequences as Probes

5           The identified sequence of the present invention can be used as probes to obtain the sequence of additional genes of interest from a second organism. For example, probes to potential bacterial target proteins may be hybridized to nucleic acids from other organisms including other bacteria and higher organisms, to identify homologous sequences. Such hybridization might indicate that the protein encoded by the gene to which the probe corresponds is found in humans and therefore not necessarily a good drug target. Alternatively, the gene can be conserved only in bacteria and therefore would be a good drug target for a broad spectrum antibiotic or antimicrobial.

10           Probes derived from the identified nucleic acid sequences of interest or portions thereof can be labeled with detectable labels familiar to those skilled in the art, including radioisotopes and non-radioactive labels, to provide a detectable probe. The detectable probe can be single stranded or double stranded and can be made using techniques known in the art, including *in vitro* transcription, nick translation, or kinase reactions. A nucleic acid sample containing a sequence capable of hybridizing to the labeled probe is contacted with the labeled probe. If the nucleic acid in the sample is double stranded, it can be denatured prior to contacting the probe. In some applications, the nucleic acid sample can be immobilized on a surface such as a nitrocellulose or nylon membrane. The nucleic acid sample can comprise nucleic acids obtained from a variety of sources, including genomic DNA, cDNA libraries, RNA, or tissue samples.

Procedures used to detect the presence of nucleic acids capable of hybridizing to the detectable probe include well known techniques such as Southern blotting, Northern blotting, dot blotting, colony hybridization, and plaque hybridization. In some applications, the nucleic acid capable of hybridizing to the labeled probe can be cloned into vectors such as expression vectors, sequencing vectors, or in vitro transcription vectors to facilitate the characterization and expression of the hybridizing nucleic acids in the sample. For example, such techniques can be used to isolate, purify and clone sequences from a genomic library, made from a variety of bacterial species, which are capable of hybridizing to probes made from the sequences identified in Examples 5 and 6.

#### EXAMPLE 14

##### Preparation of PCR Primers and Amplification of DNA

The identified *E. coli* genes corresponding directly to or located within the operon of nucleic acid sequences required for proliferation or portions thereof can be used to prepare PCR primers for a variety of applications, including the identification or isolation of homologous sequences from other species, for example *S. typhimurium*, *E. cloacae*, and *Klebsiella pneumoniae*, which contain part or all of the homologous genes. Because homologous genes are related but not identical in sequence, those skilled in the art will often employ degenerate sequence PCR primers. Such degenerate sequence primers are designed based on conserved sequence regions, either known or suspected, such as conserved coding regions. The successful production of a PCR product using degenerate probes generated from the sequences identified herein would indicate the presence of a homologous gene sequence in the species being screened. The PCR primers are at least 10 bases, and preferably at least 20 bases in length. More preferably, the PCR primers are at least 20-30 bases in length. In some embodiments, the PCR primers can be more than 30 bases in length. It is preferred that the primer pairs have approximately the same G/C ratio, so that melting temperatures are approximately the same. A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR technology, see Molecular Cloning to Genetic Engineering White, B.A. Ed. in **Methods in Molecular Biology** 67: Humana Press, Totowa 1997. When the entire coding sequence of the target gene is known, the 5' and 3' regions of the target gene can be used as the sequence source for PCR probe generation. In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid

sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites.

### EXAMPLE 15

#### Inverse PCR

The technique of inverse polymerase chain reaction can be used to extend the known nucleic acid sequence identified in Examples 5 and 6. The inverse PCR reaction is described generally by Ochman et al., in Ch. 10 of **PCR Technology: Principles and Applications for DNA Amplification**, (Henry A. Erlich, Ed.) W.H. Freeman and Co. (1992). Traditional PCR requires two primers that are used to prime the synthesis of complementary strands of DNA. In inverse PCR, only a core sequence need be known.

Using the sequences identified as relevant from the techniques taught in Examples 5 and 6 and applied to other species of bacteria, a subset of exogenous nucleic sequences are identified that correspond to genes or operons that are required for bacterial proliferation. In species for which a genome sequence is not known, the technique of inverse PCR provides a method for obtaining the gene in order to determine the sequence or to place the probe sequences in full context to the target sequence to which the identified exogenous nucleic acid sequence binds.

To practice this technique, the genome of the target organism is digested with an appropriate restriction enzyme so as to create fragments of nucleic acid that contain the identified sequence as well as unknown sequences that flank the identified sequence. These fragments are then circularized and become the template for the PCR reaction. PCR primers are designed in accordance with the teachings of Example 15 and directed to the ends of the identified sequence are synthesized. The primers direct nucleic acid synthesis away from the known sequence and toward the unknown sequence contained within the circularized template. After the PCR reaction is complete, the resulting PCR products can be sequenced so as to extend the sequence of the identified gene past the core sequence of the identified exogenous nucleic acid sequence identified. In this manner, the full

sequence of each novel gene can be identified. Additionally the sequences of adjacent coding and noncoding regions can be identified.

#### EXAMPLE 16

##### Identification of Genes Required for *Staphylococcus aureus* Proliferation

Genes required for proliferation in *Staphylococcus aureus* are identified according to the methods described above.

#### EXAMPLE 17

##### Identification of Genes Required for *Neisseria gonorrhoeae* Proliferation

Genes required for proliferation in *Neisseria gonorrhoeae* are identified according to the methods described above.

#### EXAMPLE 18

##### Identification of Genes Required for *Pseudomonas aeruginosa* Proliferation

Genes required for proliferation in *Pseudomonas aeruginosa* are identified according to the methods described above.

#### EXAMPLE 19

##### Identification of Genes Required for *Enterococcus faecalis* Proliferation

Genes required for proliferation in *Enterococcus faecalis* are identified according to the methods described above.

#### EXAMPLE 20

##### Identification of Genes Required for *Haemophilus influenzae* Proliferation

Genes required for proliferation in *Haemophilus influenzae* are identified according to the methods described above.

#### EXAMPLE 21

##### Identification of Genes Required for *Salmonella typhimurium* Proliferation

Genes required for proliferation in *Salmonella typhimurium* are identified according to the methods described above.

#### EXAMPLE 22

##### Identification of Genes Required for *Helicobacter pylori* Proliferation

Genes required for proliferation in *Helicobacter pylori* are identified according to the methods described above.

#### EXAMPLE 23

##### Identification of Genes Required for *Mycoplasma pneumoniae* Proliferation

Genes required for proliferation in *Mycoplasma pneumoniae* are identified according to the methods described above.

#### EXAMPLE 24

##### Identification of Genes Required for *Plasmodium ovale* Proliferation

5 Genes required for proliferation in *Plasmodium ovale* are identified according to the methods described above.

#### EXAMPLE 25

##### Identification of Genes Required for *Saccharomyces cerevisiae* Proliferation

10 Genes required for proliferation in *Saccharomyces cerevisiae* are identified according to the methods described above.

#### EXAMPLE 26

##### Identification of Genes Required for *Entamoeba histolytica* Proliferation

15 Genes required for proliferation in *Entamoeba histolytica* are identified according to the methods described above.

#### EXAMPLE 27

##### Identification of Genes Required for *Candida albicans* Proliferation

Genes required for proliferation in *Candida albicans* are identified according to the methods described above.

#### EXAMPLE 28

##### Identification of Genes Required for *Klebsiella pneumoniae* Proliferation

20 Genes required for proliferation in *Klebsiella pneumoniae* are identified according to the methods described above.

#### EXAMPLE 29

##### Identification of Genes Required for *Salmonella typhi* Proliferation

25 Genes required for proliferation in *Salmonella typhi* are identified according to the methods described above.

#### EXAMPLE 30

##### Identification of Genes Required for *Salmonella paratyphi* Proliferation

30 Genes required for proliferation in *Salmonella paratyphi* are identified according to the methods described above.

#### EXAMPLE 31

##### Identification of Genes Required for *Salmonella choleraesuis* Proliferation



Genes required for proliferation in *Salmonella choleraesuis* are identified according to the methods described above.

#### EXAMPLE 32

##### Identification of Genes Required for *Staphylococcus epidermis* Proliferation

5 Genes required for proliferation in *Staphylococcus epidermis* are identified according to the methods described above.

#### EXAMPLE 33

##### Identification of Genes Required for *Mycobacterium tuberculosis* Proliferation

10 Genes required for proliferation in *Mycobacterium tuberculosis* are identified according to the methods described above.

#### EXAMPLE 34

##### Identification of Genes Required for *Mycobacterium leprae* Proliferation

Genes required for proliferation in *Mycobacterium leprae* are identified according to the methods described above.

15

#### EXAMPLE 35

##### Identification of Genes Required for *Treponema pallidum* Proliferation

Genes required for proliferation in *Treponema pallidum* are identified according to the methods described above.

#### EXAMPLE 36

20

##### Identification of Genes Required for *Bacillus anthracis* Proliferation

Genes required for proliferation in *Bacillus anthracis* are identified according to the methods described above.

#### EXAMPLE 37

##### Identification of Genes Required for *Yersinia pestis* Proliferation

25

Genes required for proliferation in *Yersinia pestis* are identified according to the methods described above.

#### EXAMPLE 38

##### Identification of Genes Required for *Clostridium botulinum* Proliferation

30 Genes required for proliferation in *Clostridium botulinum* are identified according to the methods described above.

#### EXAMPLE 39

##### Identification of Genes Required for *Campylobacter jejuni* Proliferation

Genes required for proliferation in *Campylobacter jejuni* are identified according to the methods described above.

#### EXAMPLE 40

##### Identification of Genes Required for *Chlamydia trachomatis* Proliferation

5 Genes required for proliferation in *Chlamydia trachomatis* are identified according to the methods described above.

##### **Use of Isolated Exogenous Nucleic Acid Fragments as Antisense Antibiotics**

10 In addition to using the identified sequences to enable screening of molecule libraries to identify compounds useful to identify antibiotics, the sequences themselves can be used as therapeutic agents. Specifically, the identified exogenous sequences in an antisense orientation can be provided to an individual to inhibit the translation of a bacterial target gene.

##### Generation of Antisense Therapeutics from Identified Exogenous Sequences

15 The sequences of the present invention can be used as antisense therapeutics for the treatment of bacterial infections or simply for inhibition of bacterial growth *in vitro* or *in vivo*. The therapy exploits the biological process in cells where genes are transcribed into messenger RNA (mRNA) that is then translated into proteins. Antisense RNA technology contemplates the use of antisense oligonucleotides directed  
20 against a target gene that will bind to its target and decrease or inhibit the translation of the target mRNA. In one embodiment, antisense oligonucleotides can be used to treat and control a bacterial infection of a cell culture containing a population of desired cells contaminated with bacteria. In another embodiment, the antisense oligonucleotides can be used to treat an organism with a bacterial infection.

25 Antisense oligonucleotides can be synthesized from any of the sequences of the present invention using methods well known in the art. In a preferred embodiment, antisense oligonucleotides are synthesized using artificial means. Uhlmann & Peymann, Chemical Rev. 90:543-584 (1990) review antisense oligonucleotide technology in detail. Modified or unmodified antisense oligonucleotides can be used as  
30 therapeutic agents. Modified antisense oligonucleotides are preferred since it is well known that antisense oligonucleotides are extremely unstable. Modification of the phosphate backbones of the antisense oligonucleotides can be achieved by substituting

the internucleotide phosphate residues with methylphosphonates, phosphorothioates, phosphoramidates, and phosphate esters. Nonphosphate internucleotide analogs such as siloxane bridges, carbonate bridges, thioester bridges, as well as many others known in the art. The preparation of certain antisense oligonucleotides with modified internucleotide linkages is described in U.S. Patent No. 5,142,047, hereby incorporated by reference.

Modifications to the nucleoside units of the antisense oligonucleotides are also contemplated. These modifications can increase the half-life and increase cellular rates of uptake for the oligonucleotides *in vivo*. For example,  $\alpha$ -anomeric nucleotide units and modified bases such as 1,2-dideoxy-d-ribofuranose, 1,2-dideoxy-1-phenylribofuranose, and  $N^4$ ,  $N^4$ -ethano-5-methyl-cytosine are contemplated for use in the present invention.

An additional form of modified antisense molecules is found in peptide nucleic acids. Peptide nucleic acids (PNA) have been developed to hybridize to single and double stranded nucleic acids. PNA are nucleic acid analogs in which the entire deoxyribose-phosphate backbone has been exchanged with a chemically completely different, but structurally homologous, polyamide (peptide) backbone containing 2-aminoethyl glycine units. Unlike DNA, which is highly negatively charged, the PNA backbone is neutral. Therefore, there is much less repulsive energy between complementary strands in a PNA-DNA hybrid than in the comparable DNA-DNA hybrid, and consequently they are much more stable. PNA can hybridize to DNA in either a Watson/Crick or Hoogsteen fashion (Demidov et al., *Proc. Natl. Acad. Sci. U.S.A.* **92**:2637-2641, 1995; Egholm, *Nature* **365**:566-568, 1993; Nielsen et al., *Science* **254**:1497-1500, 1991; Dueholm et al., *New J. Chem.* **21**:19-31, 1997).

Molecules called PNA "clamps" have been synthesized which have two identical PNA sequences joined by a flexible hairpin linker containing three 8-amino-3,6-dioxaoctanoic acid units. When a PNA clamp is mixed with a complementary homopurine or homopyrimidine DNA target sequence, a PNA-DNA-PNA triplex hybrid can form which has been shown to be extremely stable (Bentin et al., *Biochemistry* **35**:8863-8869, 1996; Egholm et al., *Nucleic Acids Res.* **23**:217-222, 1995; Griffith et al., *J. Am. Chem. Soc.* **117**:831-832, 1995).

The sequence-specific and high affinity duplex and triplex binding of PNA have been extensively described (Nielsen et al., *Science* **254**:1497-1500, 1991; Egholm et al., *J. Am. Chem. Soc.* **114**:9677-9678, 1992; Egholm et al., *Nature* **365**:566-568, 1993; Almarsson et al., *Proc. Natl. Acad. Sci. U.S.A.* **90**:9542-9546, 1993; Demidov et al., *Proc. Natl. Acad. Sci. U.S.A.* **92**:2637-2641, 1995). They have also been shown to be resistant to nuclease and protease digestion (Demidov et al., *Biochem. Pharm.* **48**:1010-1313, 1994). PNA has been used to inhibit gene expression (Hanvey et al., *Science* **258**:1481-1485, 1992; Nielsen et al., *Nucl. Acids. Res.*, **21**:197-200, 1993; Nielsen et al., *Gene* **149**:139-145, 1994; Good & Nielsen, *Science*, **95**: 2073-2076, 1998; all of which are hereby incorporated by reference), to block restriction enzyme activity (Nielsen et al., *supra.*, 1993), to act as an artificial transcription promoter (Mollegaard, *Proc. Natl. Acad. Sci. U.S.A.* **91**:3892-3895, 1994) and as a pseudo restriction endonuclease (Demidov et al., *Nucl. Acids. Res.* **21**:2103-2107, 1993). Recently, PNA has also been shown to have antiviral and antitumoral activity mediated through an antisense mechanism (Norton, *Nature Biotechnol.*, **14**:615-619, 1996; Hirschman et al., *J. Investig. Med.* **44**:347-351, 1996). PNAs have been linked to various peptides in order to promote PNA entry into cells (Basu et al., *Bioconj. Chem.* **8**:481-488, 1997; Pardridge et al., *Proc. Natl. Acad. Sci. U.S.A.* **92**:5592-5596, 1995).

The antisense oligonucleotides contemplated by the present invention can be administered by direct application of oligonucleotides to a target using standard techniques well known in the art. The antisense oligonucleotides can be generated within the target using a plasmid, or a phage. Alternatively, the antisense nucleic acid may be expressed from a sequence in the chromosome of the target cell. It is further contemplated that the antisense oligonucleotide contemplated are incorporated in a ribozyme sequence to enable the antisense to specifically bind and cleave its target mRNA. For technical applications of ribozyme and antisense oligonucleotides see Rossi et al., *Pharmacol. Ther.* **50**(2):245-254, (1991), which is hereby incorporated by reference. The present invention also contemplates using a retron to introduce an antisense oligonucleotide to a cell. Retron technology is exemplified by U.S. Patent No. 5,405,775, which is hereby incorporated by reference. Antisense oligonucleotides can also be delivered using liposomes or by electroporation techniques which are well known in the art.

The antisense nucleic acids of the present invention can also be used to design antibiotic compounds comprising nucleic acids which function by intracellular triple helix formation. Triple helix oligonucleotides are used to inhibit transcription from a genome. The sequences identified as required for proliferation in the present invention, or portions thereof, can be used as templates to inhibit microorganism gene expression in individuals infected with such organisms. Traditionally, homopurine sequences were considered the most useful for triple helix strategies. However, homopyrimidine sequences can also inhibit gene expression. Such homopyrimidine oligonucleotides bind to the major groove at homopurine:homopyrimidine sequences. Thus, both types of sequences based on the sequences of the present invention that are required for proliferation are contemplated for use as antibiotic compound templates.

The antisense oligonucleotides of this example employ the identified sequences of the present invention to induce bacterial cell death or at least bacterial stasis by inhibiting target gene translation. Antisense oligonucleotides containing from about 8 to 40 bases of the sequences of the present invention have sufficient complementarity to form a duplex with the target sequence under physiological conditions.

To kill bacterial cells or inhibit their growth, the antisense oligonucleotides are applied to the bacteria or to the target cells under conditions that facilitate their uptake. These conditions include sufficient incubation times of cells and oligonucleotides so that the antisense oligonucleotides are taken up by the cells. In one embodiment, an incubation period of 7-10 days is sufficient to kill bacteria in a sample. An optimum concentration of antisense oligonucleotides is selected for use.

The concentration of antisense oligonucleotides to be used can vary depending on the type of bacteria sought to be controlled, the nature of the antisense oligonucleotide to be used, and the relative toxicity of the antisense oligonucleotide to the desired cells in the treated culture. Antisense oligonucleotides can be introduced to cell samples at a number of different concentrations preferably between  $1 \times 10^{-10}$  M to  $1 \times 10^{-4}$  M. Once the minimum concentration that can adequately control gene expression is identified, the optimized dose is translated into a dosage suitable for use *in vivo*. For example, an inhibiting concentration in culture of  $1 \times 10^{-7}$  translates into a dose of approximately 0.6 mg/kg body weight. Levels of oligonucleotide approaching 100 mg/kg body weight or higher may be possible after testing the toxicity of the oligonucleotide in

laboratory animals. It is additionally contemplated that cells from the subject are removed, treated with the antisense oligonucleotide, and reintroduced into the subject. This range is merely illustrative and one of skill in the art are able to determine the optimal concentration to be used in a given case.

5           After the bacterial cells have been killed or controlled in a desired culture, the desired cell population may be used for other purposes.

#### EXAMPLE 41

10           The following example demonstrates the ability of an *E. coli* antisense oligonucleotide to act as a bactericidal or bacteriostatic agent to treat a contaminated cell culture system. The application of the antisense oligonucleotides of the present invention are thought to inhibit the translation of bacterial gene products required for proliferation.

15           The antisense oligonucleotide of this example corresponds to a 30 base phosphorothioate modified oligodeoxynucleotide complementary to a nucleic acid involved in proliferation, such as Molecule Number EcXA001. A sense oligodeoxynucleotide complementary to the antisense sequence is synthesized and used as a control. The oligonucleotides are synthesized and purified according to the procedures of Matsukura, et al., Gene 72:343 (1988). The test oligonucleotides are dissolved in a small volume of autoclaved water and added to culture medium to make a  
20           100 micromolar stock solution.

          Human bone marrow cells are obtained from the peripheral blood of two patients and cultured according standard procedures well known in the art. The culture is contaminated with the K-12 strain of *E. coli* and incubated at 37°C overnight to establish bacterial infection.

25           The control and antisense oligonucleotide containing solutions are added to the contaminated cultures and monitored for bacterial growth. After a 10 hour incubation of culture and oligonucleotides, samples from the control and experimental cultures are drawn and analyzed for the translation of the target bacterial gene using standard microbiological techniques well known in the art. The target *E. coli* gene is found to be  
30           translated in the control culture treated with the control oligonucleotide, however, translation of the target gene in the experimental culture treated with the antisense oligonucleotide of the present invention is not detected or reduced.

#### EXAMPLE 42

A subject suffering from an *E. coli* infection is treated with the antisense oligonucleotide preparation of Example 39. The antisense oligonucleotide is provided in a pharmaceutically acceptable carrier at a concentration effective to inhibit the translation of the target gene. The present subject is treated with a concentration of antisense oligonucleotide sufficient to achieve a blood concentration of about 100 micromolar. The patient receives daily injections of antisense oligonucleotide to maintain this concentration for a period of 1 week. At the end of the week a blood sample is drawn and analyzed for the presence or absence using standard techniques well known in the art. There is no detectable evidence of *E. coli* and the treatment is terminated.

#### EXAMPLE 43

##### Preparation and use of Triple Helix Probes

The sequences of microorganism genes required for proliferation of the present invention are scanned to identify 10-mer to 20-mer homopyrimidine or homopurine stretches that could be used in triple-helix based strategies for inhibiting gene expression. Following identification of candidate homopyrimidine or homopurine stretches, their efficiency in inhibiting gene expression is assessed by introducing varying amounts of oligonucleotides containing the candidate sequences into a population of bacterial cells that normally express the target gene. The oligonucleotides may be prepared on an oligonucleotide synthesizer or they may be purchased commercially from a company specializing in custom oligonucleotide synthesis, such as GENSET, Paris, France.

The oligonucleotides can be introduced into the cells using a variety of methods known to those skilled in the art, including but not limited to calcium phosphate precipitation, DEAE-Dextran, electroporation, liposome-mediated transfection or native uptake.

Treated cells are monitored for a reduction in proliferation using techniques such as monitoring growth levels as compared to untreated cells using optical density measurements. The oligonucleotides that are effective in inhibiting gene expression in cultured cells can then be introduced *in vivo* using the techniques well known in that art at a dosage level shown to be effective.

In some embodiments, the natural (beta) anomers of the oligonucleotide units can be replaced with alpha anomers to render the oligonucleotide more resistant to nucleases. Further, an intercalating agent such as ethidium bromide, or the like, can be attached to the 3' end of the alpha oligonucleotide to stabilize the triple helix. For information on the generation of oligonucleotides suitable for triple helix formation see Griffin et al. (**Science** 245:967-971 (1989), which is hereby incorporated by this reference).

#### **EXAMPLE 44**

##### Identification of Bacterial Strains from Isolated Specimens by PCR

Classical bacteriological methods for the detection of various bacterial species are time consuming and costly. These methods include growing the bacteria isolated from a subject in specialized media, cultivation on selective agar media, followed by a set of confirmation assays that can take from 8 to 10 days or longer to complete. Use of the identified sequences of the present invention provides a method to dramatically reduce the time necessary to detect and identify specific bacterial species present in a sample.

In one exemplary method, bacteria are grown in enriched media and DNA samples are isolated from specimens of, for example, blood, urine, stool, saliva or central nervous system fluid by conventional methods. A panel of PCR primers based on identified sequences unique to various species of microorganisms are then utilized in accordance with Example 12 to amplify DNA of approximately 100-200 bases in length from the specimen. A separate PCR reaction is set up for each pair of PCR primers and after the PCR reaction is complete, the reaction mixtures are assayed for the presence of PCR product. The presence or absence of bacteria from the species to which the PCR primer pairs belong is determined by the presence or absence of a PCR product in the various test PCR reaction tubes.

Although the PCR reaction is used to assay the isolated sample for the presence of various bacterial species, other assays such as the Southern blot hybridization are also contemplated.

All documents cited herein are incorporated herein by reference in their entireties.